


AQP3 Polyclonal Antibody

Catalog No :	YT0291
Reactivity :	Human;Mouse;Rat
Applications :	IHC;IF;ELISA
Target :	AQP3
Fields :	>>Vasopressin-regulated water reabsorption
Gene Name :	AQP3
Protein Name :	Aquaporin-3
Human Gene Id :	360
Human Swiss Prot No :	Q92482
Mouse Gene Id :	11828
Mouse Swiss Prot No :	Q8R2N1
Rat Gene Id :	65133
Rat Swiss Prot No :	P47862
Immunogen :	The antiserum was produced against synthesized peptide derived from human AQP3. AA range:165-214
Specificity :	AQP3 Polyclonal Antibody detects endogenous levels of AQP3 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	32kD
Background :	This gene encodes the water channel protein aquaporin 3. Aquaporins are a family of small integral membrane proteins related to the major intrinsic protein, also known as aquaporin 0. Aquaporin 3 is localized at the basal lateral membranes of collecting duct cells in the kidney. In addition to its water channel function, aquaporin 3 has been found to facilitate the transport of nonionic small solutes such as urea and glycerol, but to a smaller degree. It has been suggested that water channels can be functionally heterogeneous and possess water and solute permeation mechanisms. Alternative splicing of this gene results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Dec 2015],
Function :	domain:Aquaporins contain two tandem repeats each containing three membrane-spanning domains and a pore-forming loop with the signature motif Asn-Pro-Ala (NPA).,function:Forms a water-specific channel that provide kidney medullary collecting duct with high permeability to water, thereby permitting water to move in the direction of an osmotic gradient. May function as a water and urea exit mechanism in antidiuresis in collecting duct cells. Also slightly permeable to urea and glycerol. May play an important role in gastrointestinal tract water transport and in glycerol metabolism.,online information:Blood group antigen gene mutation database,polymorphism:AQP3 is responsible for the GIL blood group system. Isoform 2 is detected in GIL-negative individuals that lack functional AQP3.,similarity:Belongs to the MIP/aquaporin (TC 1.A.8) family.,subcellular location:In collecting ducts of kidney
Subcellular Location :	Cell membrane ; Multi-pass membrane protein . Basolateral cell membrane ; Multi-pass membrane protein .
Expression :	Widely expressed in epithelial cells of kidney (collecting ducts) and airways, in keratinocytes, immature dendritic cells and erythrocytes. Isoform 2 is not detectable in erythrocytes at the protein level.
Sort :	2183
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Immunohistochemistry analysis of AQP3 antibody in paraffin-embedded human lung carcinoma tissue.