


BCAS2 Polyclonal Antibody


| | |
|------------------------------|---|
| Catalog No : | YT0464 |
| Reactivity : | Human;Mouse;Rat;Monkey |
| Applications : | WB;ELISA |
| Target : | BCAS2 |
| Fields : | >>Spliceosome |
| Gene Name : | BCAS2 |
| Protein Name : | Pre-mRNA-splicing factor SPF27 |
| Human Gene Id : | 10286 |
| Human Swiss Prot No : | O75934 |
| Mouse Gene Id : | 68183 |
| Mouse Swiss Prot No : | Q9D287 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human BCAS2. AA range:176-225 |
| Specificity : | BCAS2 Polyclonal Antibody detects endogenous levels of BCAS2 protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | WB 1:500 - 1:2000. ELISA: 1:5000. Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |

| | |
|-------------------------------|--|
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |
| Observed Band : | 27kD |
| Cell Pathway : | Spliceosome; |
| Background : | function:Involved in mRNA splicing.,similarity:Belongs to the SPF27 family.,subunit:Associated with the spliceosome.,tissue specificity:Ubiquitously expressed., |
| Function : | function:Involved in mRNA splicing.,similarity:Belongs to the SPF27 family.,subunit:Associated with the spliceosome.,tissue specificity:Ubiquitously expressed., |
| Subcellular Location : | Nucleus . Nucleus, nucleolus . |
| Expression : | Ubiquitously expressed. |
| Sort : | 2606 |
| No4 : | 1 |
| Host : | Rabbit |
| Modifications : | Unmodified |


Products Images



Western Blot analysis of various cells using BCAS2 Polyclonal Antibody diluted at 1:1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Western Blot analysis of COS7 cells using BCAS2 Polyclonal Antibody diluted at 1:1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Western blot analysis of the lysates from HUVEC cells using BCAS2 antibody.