


Carbonyl Reductase 1 Polyclonal Antibody

Catalog No :	YT0632
Reactivity :	Human;Rat;Mouse;
Applications :	WB;IHC;IF;ELISA
Target :	Carbonyl Reductase 1
Fields :	>>Arachidonic acid metabolism;>>Folate biosynthesis;>>Metabolism of xenobiotics by cytochrome P450;>>Metabolic pathways;>>Chemical carcinogenesis - DNA adducts;>>Chemical carcinogenesis - reactive oxygen species
Gene Name :	CBR1
Protein Name :	Carbonyl reductase [NADPH] 1
Human Gene Id :	873
Human Swiss Prot No :	P16152
Mouse Swiss Prot No :	P48758
Immunogen :	The antiserum was produced against synthesized peptide derived from human CBR1. AA range:181-230
Specificity :	Carbonyl Reductase 1 Polyclonal Antibody detects endogenous levels of Carbonyl Reductase 1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.



Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	32kD
Cell Pathway :	Arachidonic acid metabolism;
Background :	The protein encoded by this gene belongs to the short-chain dehydrogenases/reductases (SDR) family, which function as NADPH-dependent oxidoreductases having wide specificity for carbonyl compounds, such as quinones, prostaglandins, and various xenobiotics. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Nov 2013],
Function :	<p>catalytic activity:(13E)-(15S)-11-alpha,15-dihydroxy-9-oxoprost-13-enoate + NADP(+) = (13E)-11-alpha-hydroxy-9,15-dioxoprost-13-enoate + NADPH.,catalytic activity:(5Z,13E)-(15S)-9-alpha,11-alpha,15-trihydroxyprosta-5,13-dienoate + NADP(+) = (5Z,13E)-(15S)-11-alpha,15-dihydroxy-9-oxoprost-5,13-dienoate + NADPH.,catalytic activity:R-CHOH-R' + NADP(+) = R-CO-R' + NADPH.,function:Catalyzes the reduction of a wide variety of carbonyl compounds including the antitumor anthracycline antibiotics. Can convert prostaglandin E2 to prostaglandin F2-alpha.,similarity:Belongs to the short-chain dehydrogenases/reductases (SDR) family.,subunit:Monomer.,</p>
Subcellular Location :	Cytoplasm .
Expression :	Expressed in kidney (at protein level).
Sort :	3113
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images

Western Blot analysis of 293T HELA using Carbonyl Reductase 1 Polyclonal Antibody. Antibody was diluted at 1:1000



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using CBR1 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa cells, using CBR1 Antibody. The lane on the right is blocked with the synthesized peptide.