

CD137 Polyclonal Antibody

YT0722 Catalog No:

Reactivity: Human; Rat; Mouse;

Applications: WB;IF;ELISA

Target: CD137

Fields: >>Cytokine-cytokine receptor interaction

Gene Name: TNFRSF9

Tumor necrosis factor receptor superfamily member 9 **Protein Name:**

Human Gene Id: 3604

Human Swiss Prot

Q07011

No:

Mouse Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

TNFRSF9. AA range:101-150

CD137 Polyclonal Antibody detects endogenous levels of CD137 protein. **Specificity:**

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other

applications.

P20334

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/3



Observed Band: 27kD

Cell Pathway: Cytokine-cytokine receptor interaction;

Background : The protein encoded by this gene is a member of the TNF-receptor superfamily.

This receptor contributes to the clonal expansion, survival, and development of T cells. It can also induce proliferation in peripheral monocytes, enhance T cell apoptosis induced by TCR/CD3 triggered activation, and regulate CD28 costimulation to promote Th1 cell responses. The expression of this receptor is induced by lymphocyte activation. TRAF adaptor proteins have been shown to bind to this receptor and transduce the signals leading to activation of NF-

kappaB. [provided by RefSeg, Jul 2008],

Function: function:Receptor for TNFSF14/4-1BBL. Possibly active during T cell

activation., similarity: Contains 4 TNFR-Cys repeats., subunit: Interacts with TRAF1,

TRAF2 and TRAF3. Interacts with LRR-repeat protein 1/LRR-1.,tissue

specificity:Expressed on the surface of activated T-cells.,

Subcellular

Location:

Membrane; Single-pass type I membrane protein.

Expression: Expressed on the surface of activated T-cells.


Sort : 3378

No4:


Host: Rabbit

Modifications: Unmodified

Products Images



Western Blot analysis of mouse-brain cells using CD137 Polyclonal Antibody diluted at 1:500



Western blot analysis of lysates from K562 cells, using TNFRSF9 Antibody. The lane on the right is blocked with the synthesized peptide.