

## AZ1 Polyclonal Antibody


<b>Catalog No :</b>	YT0427
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;IHC;IF;ELISA
<b>Target :</b>	AZ1
<b>Gene Name :</b>	OAZ1
<b>Protein Name :</b>	Ornithine decarboxylase antizyme 1
<b>Human Gene Id :</b>	4946
<b>Human Swiss Prot No :</b>	P54368
<b>Mouse Gene Id :</b>	18245
<b>Mouse Swiss Prot No :</b>	P54369
<b>Rat Gene Id :</b>	25502
<b>Rat Swiss Prot No :</b>	P54370
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human OAZ1. AA range:14-63
<b>Specificity :</b>	AZ1 Polyclonal Antibody detects endogenous levels of AZ1 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	IHC 1:100 - 1:300. ELISA: 1:40000, WB 1:500-2000. IF 1:50-200
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

---

<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)
<b>Molecularweight :</b>	25kD
<b>Background :</b>	<p>The protein encoded by this gene belongs to the ornithine decarboxylase antizyme family, which plays a role in cell growth and proliferation by regulating intracellular polyamine levels. Expression of antizymes requires +1 ribosomal frameshifting, which is enhanced by high levels of polyamines. Antizymes in turn bind to and inhibit ornithine decarboxylase (ODC), the key enzyme in polyamine biosynthesis; thus, completing the auto-regulatory circuit. This gene encodes antizyme 1, the first member of the antizyme family, that has broad tissue distribution, and negatively regulates intracellular polyamine levels by binding to and targeting ODC for degradation, as well as inhibiting polyamine uptake. Antizyme 1 mRNA contains two potential in-frame AUGs; and studies in rat suggest that alternative use of the two translation initiation sites results in N-terminally distinct protein isoforms</p>
<b>Function :</b>	<p>alternative products:A ribosomal frameshift occurs between the codons for Ser-68 and Asp-69. An autoregulatory mechanism enables modulation of frameshifting according to the cellular concentration of polyamines,function:Binds to, and destabilizes, ornithine decarboxylase which is then degraded. Also inhibits cellular uptake of polyamines by inactivating the polyamine uptake transporter.,similarity:Belongs to the ODC antizyme family.,</p>
<b>Subcellular Location :</b>	nucleus,cytoplasm,cytosol,
<b>Expression :</b>	Brain,Fibroblast,Lymphoma,
<b>Tag :</b>	orthogonal
<b>Sort :</b>	2539
<b>No4 :</b>	1
<b>Host :</b>	Rabbit
<b>Modifications :</b>	Unmodified

---

## Products Images



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using OAZ1 Antibody. The picture on the right is blocked with the synthesized peptide.