

IKK α/β (PT0435R) PT® Rabbit mAb

Catalog No: YM8275

Reactivity: Human; Mouse; Rat;

Applications: WB;IHC;IF;IP;ELISA

Target: IKKα/β

Fields: >>Antifolate resistance;>>MAPK signaling pathway;>>Ras signaling

pathway;>>Chemokine signaling pathway;>>NF-kappa B signaling

pathway;>>FoxO signaling pathway;>>mTOR signaling pathway;>>PI3K-Akt signaling pathway;>>Apoptosis;>>Osteoclast differentiation;>>Toll-like receptor signaling pathway;>>NOD-like receptor signaling pathway;>>RIG-I-like receptor signaling pathway;>>Cytosolic DNA-sensing pathway;>>C-type lectin receptor

signaling pathway;>>IL-17 signaling pathway;>>Th1 and Th2 cell

differentiation;>>Th17 cell differentiation;>>T cell receptor signaling pathway;>>B

cell receptor signaling pathway;>>TNF signaling pathway;>>Adipocytokine signaling pathway;>>Alcoholic liver disease;>>Alzheimer disease;>>Epithelial cell

signaling in Helicobacter pylori infection;>>Pathogenic Escherichia coli

infection;>>Shigellosis;>>Salmonella infection;>>Yersinia infection;>>Chagas disease;>>Toxoplasmosis;>>Hepatitis C;>>Hepatitis B;>>Measles;>>Human

cytomegalovirus infection;>>Influenza A;>>Human pap

Gene Name: CHUK/IKBKB

Protein Name: Inhibitor of nuclear factor kappa-B kinase subunit alpha

Human Gene Id: 1147/3551

Human Swiss Prot 015111:014920

No:

Mouse Gene Id: 16150

Rat Gene Id: 84351

Rat Swiss Prot No: Q9QY78

Specificity: endogenous

1/3



Formulation: PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA

Source: Monoclonal, rabbit, lgG, Kappa

Dilution: IHC 1:500-1:2000;WB 1:1000-1:5000;IF 1:200-1:1000;ELISA

1:5000-1:20000;IP 1:50-1:200;

Purification: Protein A

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 86kD

Observed Band: 86kD

Cell Pathway: T_Cell_Receptor; Insulin Receptor; B_Cell_Antigen; Stem cell pathway;

Toll Like; MAPK ERK Growth; MAPK G Protein; PI3K/Akt; NF kappaB;

Protein_Acetylation

Background: This gene encodes a member of the serine/threonine protein kinase family. The

encoded protein, a component of a cytokine-activated protein complex that is an inhibitor of the essential transcription factor NF-kappa-B complex, phosphorylates sites that trigger the degradation of the inhibitor via the ubiquination pathway, thereby activating the transcription factor. [provided by RefSeq, Jul 2008],

Function : catalytic activity:ATP + [I-kappa-B protein] = ADP + [I-kappa-B

phosphoprotein].,enzyme regulation: Activated when phosphorylated and

inactivated when dephosphorylated.,function: Acts as part of the IKK complex in the conventional pathway of NF-kappa-B activation and phosphorylates inhibitors

of NF-kappa-B thus leading to the dissociation of the inhibitor/NF-kappa-B complex and ultimately the degradation of the inhibitor. As part of the non-

canonical pathway of NF-kappa-B activation, the MAP3K14-activated CHUK/IKKA homodimer phosphorylates NFKB2/p100 associated with RelB, inducing its proteolytic processing to NFKB2/p52 and the formation of NF-kappa-B RelB-p52 complexes. Also phosphorylates NCOA3. Phosphorylates 'Ser-10' of histone H3 at NF-kappa-B-regulated promoters during inflammatory responses

triggered by cytokines., PTM: Phosphorylated by MAP3K14/NIK, AKT and to a

lesser extent by MEKK

Subcellular Location:

Cytoplasm, Nucleus

Expression: Widely expressed.

Tag: hot,recombinant

Sort: 1


No3: ab178870

No4: 1


Host: Rabbit

Modifications: Unmodified

Products Images



Various whole cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with anti-IKK α/β (PT0435R) antibody. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody. Lane 1: HEK293 Lane 2: Hela Lane 3: Rat womb Predicted band size: 86kDa Observed band size: 86kDa



Human breast carcinoma was stained with anti-IKK α/β (PT0435R) rabbit antibody