


c-FLIP Polyclonal Antibody

Catalog No :	YT0877
Reactivity :	Human;Rat;Mouse;
Applications :	WB;IHC;IF;ELISA
Target :	c-FLIP
Fields :	>>NF-kappa B signaling pathway;>>Autophagy - animal;>>Apoptosis;>>Necroptosis;>>TNF signaling pathway;>>Chagas disease;>>Hepatitis C
Gene Name :	CFLAR
Protein Name :	CASP8 and FADD-like apoptosis regulator
Human Gene Id :	8837
Human Swiss Prot No :	O15519
Mouse Swiss Prot No :	O35732
Immunogen :	The antiserum was produced against synthesized peptide derived from human in ternal CFLAR. AA range:181-230
Specificity :	c-FLIP Polyclonal Antibody detects endogenous levels of c-FLIP protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml

Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	55kD
Cell Pathway :	Apoptosis_Inhibition;Apoptosis_Mitochondrial;Apoptosis_Overview;
Background :	The protein encoded by this gene is a regulator of apoptosis and is structurally similar to caspase-8. However, the encoded protein lacks caspase activity and appears to be itself cleaved into two peptides by caspase-8. Several transcript variants encoding different isoforms have been found for this gene, and partial evidence for several more variants exists. [provided by RefSeq, Feb 2011],
Function :	domain:The caspase domain lacks the active sites residues involved in catalysis.,function:Apoptosis regulator protein which may function as a crucial link between cell survival and cell death pathways in mammalian cells. Acts as an inhibitor of TNFRSF6 mediated apoptosis. A proteolytic fragment (p43) is likely retained in the death-inducing signaling complex (DISC) thereby blocking further recruitment and processing of caspase-8 at the complex. Full length and shorter isoforms have been shown either to induce apoptosis or to reduce TNFRSF-triggered apoptosis. Lacks enzymatic (caspase) activity.,induction:Repressed by IL-2 after TCR stimulation, during progression to the S-phase of the cell cycle.,PTM:Proteolytically processed; probably by caspase-8. Processing likely occurs at the DISC and generates subunit p43 and p12.,similarity:Belongs to the peptidase C14A family.,similarity:Contains
Subcellular Location :	cytoplasm,cytosol,death-inducing signaling complex,CD95 death-inducing signaling complex,membrane raft,rioptosome,
Expression :	Widely expressed. Higher expression in skeletal muscle, pancreas, heart, kidney, placenta, and peripheral blood leukocytes. Also detected in diverse cell lines. Isoform 8 is predominantly expressed in testis and skeletal muscle.
Sort :	3884
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4 ° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).