

CD314 Polyclonal Antibody

Catalog No :	YT0753
Reactivity :	Human;Rat;Mouse;
Applications :	WB;ELISA
Target :	CD314
Fields :	>>Natural killer cell mediated cytotoxicity;>>Malaria
Gene Name :	KLRK1
Protein Name :	NKG2-D type II integral membrane protein
Human Gene Id :	22914
Human Swiss Prot No :	P26718
Mouse Swiss Prot No :	O54709
Immunogen :	The antiserum was produced against synthesized peptide derived from human KLRK1. AA range:111-160
Specificity :	CD314 Polyclonal Antibody detects endogenous levels of CD314 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 25kD

Cell Pathway : Natural killer cell mediated cytotoxicity;

Background : Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. The NKG2 gene family is located within the NK complex, a region that contains several C-type lectin genes preferentially expressed in NK cells. This gene encodes a member of the NKG2 family. The encoded transmembrane protein is characterized by a type II membrane orientation (has an extracellular C terminus) and the presence of a C-type lectin domain. It binds to a diverse family of ligands that include MHC class I chain-related A and B proteins and UL-16 binding proteins, where ligand-receptor interactions can result in the activation of

Function : alternative products:A number of isoforms are produced,function:Receptor for MICA, MICB, ULBP1, ULBP2, ULBP3 (ULBP2>ULBP1>ULBP3) and ULBP4. Plays a role as a receptor for the recognition of MHC class I HLA-E molecules by NK cells and some cytotoxic T-cells. Involved in the immune surveillance exerted by T- and B-lymphocytes.,miscellaneous:Structurally distinct families of ligands for mouse and human NKG2D receptors have been characterized. They might be orthologs.,online information:NKG-2D,similarity:Contains 1 C-type lectin domain.,subunit:Homodimer. Interacts with DAP10. The interaction with DAP10 is required for NKG2D cell surface expression.,tissue specificity:Natural killer cells. Expressed on essentially all CD56+CD3- NK cells from freshly isolated PBMC. Also detected in gamma-delta cells and CD8+ alpha-beta T-cells. Expressed in interferon-producing killer dendritic cells (IKDCs).

Subcellular Location : Cell membrane ; Single-pass type II membrane protein . Colocalized with HCST on the cell surface.

Expression : Expressed in natural killer (NK) cells, CD8(+) alpha-beta and gamma-delta T-cells. Expressed on essentially all CD56+CD3- NK cells from freshly isolated PBMC. Expressed in interferon-producing killer dendritic cells (IKDCs).


Sort : 3539

No4 : 1


Host : Rabbit

Modifications : Unmodified

Products Images



Western Blot analysis of 3T3 cells using CD314 Polyclonal Antibody



Western blot analysis of KLRK1 Antibody. The lane on the right is blocked with the KLRK1 peptide.