

CYP4A11/22 Polyclonal Antibody

Catalog No :	YT1232
Reactivity :	Human;Rat;Mouse;
Applications :	WB;ELISA
Target :	CYP4A11/22
Gene Name :	CYP4A11/CYP4A22
Protein Name :	Cytochrome P450 4A11/22
Human Gene Id :	1579/284541
Human Swiss Prot No :	Q02928/Q5TCH4
Immunogen :	The antiserum was produced against synthesized peptide derived from human Cytochrome P450 4A11/22. AA range:391-440
Specificity :	CYP4A11/22 Polyclonal Antibody detects endogenous levels of CYP4A11/22 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	60kD
Cell Pathway :	Fatty acid metabolism;Arachidonic acid metabolism;Retinol

metabolism;PPAR;Vascular smooth muscle contraction;

Background :

cytochrome P450 family 4 subfamily A member 11(CYP4A11) Homo sapiens
This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and hydroxylates medium-chain fatty acids such as laurate and myristate. Multiple transcript variants have been found for this gene. [provided by RefSeq, Jan 2016],

Function :

catalytic activity:Octane + reduced rubredoxin + O(2) = 1-octanol + oxidized rubredoxin + H(2)O.,cofactor:Heme group.,function:Catalyzes the omega- and (omega-1)-hydroxylation of various fatty acids such as laurate, myristate and palmitate. Has little activity toward prostaglandins A1 and E1. Oxidizes arachidonic acid to 20-hydroxyeicosatetraenoic acid (20-HETE).,online information:CYP4A11 alleles,polymorphism:CYP4A11v seems to be a rare allelic variant of CYP4A11, it seems to be unstable and not to metabolize lauric acid.,similarity:Belongs to the cytochrome P450 family.,tissue specificity:Kidney and liver.,

Subcellular Location :

Endoplasmic reticulum membrane ; Peripheral membrane protein . Microsome membrane ; Peripheral membrane protein .

Expression :

Expressed in liver (PubMed:7679927). Expressed in S2 and S3 segments of proximal tubules in cortex and outer medulla of kidney (PubMed:7679927, PubMed:10660572).

Sort :

4813

No4 :

1


Host :

Rabbit

Modifications :

Unmodified

Products Images



Western Blot analysis of various cells using CYP4A11/22 Polyclonal Antibody