

COL19A1 Polyclonal Antibody

Catalog No :	YT1017
Reactivity :	Human;Mouse
Applications :	WB;IHC;IF;ELISA
Target :	COL19A1
Fields :	>>Protein digestion and absorption
Gene Name :	COL19A1
Protein Name :	Collagen alpha-1(XIX) chain
Human Gene Id :	1310
Human Swiss Prot No :	Q14993
Mouse Swiss Prot No :	Q0VF58
Immunogen :	The antiserum was produced against synthesized peptide derived from human Collagen XIX alpha1. AA range:421-470
Specificity :	COL19A1 Polyclonal Antibody detects endogenous levels of COL19A1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 120kD

Background : This gene encodes the alpha chain of type XIX collagen, a member of the FACIT collagen family (fibril-associated collagens with interrupted helices). Although the function of this collagen is not known, other members of this collagen family are found in association with fibril-forming collagens such as type I and II, and serve to maintain the integrity of the extracellular matrix. The transcript produced from this gene has an unusually large 3' UTR which has not been completely sequenced. [provided by RefSeq, Jul 2008],

Function : domain:The numerous interruptions in the triple helix may make this molecule either elastic or flexible.,function:May act as a cross-bridge between fibrils and other extracellular matrix molecules. Involved in skeletal myogenesis in the developing esophagus. May play a role in organization of the pericellular matrix or the sphinteric smooth muscle.,PTM:Prolines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains.,similarity:Belongs to the fibril-associated collagens with interrupted helices (FACIT) family.,similarity:Contains 1 TSP N-terminal (TSPN) domain.,subunit:Oligomer; disulfide-linked.,tissue specificity:Localized to vascular, neuronal, mesenchymal, and some epithelial basement membrane zones in umbilical cord.,

Subcellular Location : Secreted, extracellular space, extracellular matrix .

Expression : Localized to vascular, neuronal, mesenchymal, and some epithelial basement membrane zones in umbilical cord.


Sort : 4379

No4 : 1


Host : Rabbit

Modifications : Unmodified

Products Images



Immunofluorescence analysis of A549 cells, using Collagen XIX alpha1 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human thymus gland tissue, using Collagen XIX alpha1 Antibody. The picture on the right is blocked with the synthesized peptide.