

DBC-1 Polyclonal Antibody

YT1294 Catalog No:

Reactivity: Human; Rat; Mouse;

Applications: WB;IHC;IF;ELISA

Target: DBC1

Gene Name: **KIAA1967**

Protein Name: DBIRD complex subunit KIAA1967

Q8N163

Q8VDP4

Human Gene Id: 57805

Human Swiss Prot

No:

Mouse Swiss Prot

No:

The antiserum was produced against synthesized peptide derived from human Immunogen:

KIAA1967. AA range:431-480

DBC-1 Polyclonal Antibody detects endogenous levels of DBC-1 protein. **Specificity:**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Formulation:

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. ELISA: 1:20000.. IF 1:50-200

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

-15°C to -25°C/1 year(Do not lower than -25°C) **Storage Stability:**

Observed Band: 102kD

1/3



tissue specificity: Expressed ubiquitously in normal tissues. Expressed in 84 to **Background:**

100% of neoplastic breast, lung, and colon tissues.,

Function: tissue specificity: Expressed ubiquitously in normal tissues. Expressed in 84 to

100% of neoplastic breast, lung, and colon tissues.,

Subcellular Nucleus . Cytoplasm . Cytoplasm, cytoskeleton, spindle . Recruited to chromatin, Location:

post-UV irradiation. Sequestered to the cytoplasm in the presence of MCC.

Translocated to the cytoplasm during UV-induced apoptosis. .

Expressed in gastric carcinoma tissue and the expression gradually increases **Expression:**

> with the progression of the carcinoma (at protein level). Expressed ubiquitously in normal tissues. Expressed in 84 to 100% of neoplastic breast, lung, and colon

tissues.


Sort: 5018

No4:

Host: Rabbit


Modifications: Unmodified

Products Images



Western blot analysis of lysates from Jurkat cells, using KIAA1967 Antibody. The lane on the right is blocked with the synthesized peptide.





Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).