


COL9A3 Polyclonal Antibody


Catalog No :	YT1039
Reactivity :	Human
Applications :	IHC;IF;ELISA
Target :	COL9A3
Fields :	>>PI3K-Akt signaling pathway;>>Focal adhesion;>>ECM-receptor interaction;>>Protein digestion and absorption;>>Human papillomavirus infection
Gene Name :	COL9A3
Protein Name :	Collagen alpha-3(IX) chain
Human Gene Id :	1299
Human Swiss Prot No :	Q14050
Immunogen :	The antiserum was produced against synthesized peptide derived from human Collagen IX alpha3. AA range:321-370
Specificity :	COL9A3 Polyclonal Antibody detects endogenous levels of COL9A3 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight :	64kD
Background :	This gene encodes one of the three alpha chains of type IX collagen, the major collagen component of hyaline cartilage. Type IX collagen, a heterotrimeric molecule, is usually found in tissues containing type II collagen, a fibrillar collagen. Mutations in this gene are associated with multiple epiphyseal dysplasia type 3. [provided by RefSeq, Jan 2010],
Function :	disease:Defects in COL9A3 are the cause of multiple epiphyseal dysplasia type 3 (EDM3) [MIM:600969]; also known as multiple epiphyseal dysplasia with myopathy. EDM is a generalized skeletal dysplasia associated with significant morbidity. Joint pain, joint deformity, waddling gait, and short stature are the main clinical signs and symptoms. EDM is broadly categorized into the more severe Fairbank and the milder Ribbing types.,function:Structural component of hyaline cartilage and vitreous of the eye.,PTM:Covalently linked to the telopeptides of type II collagen by lysine-derived cross-links.,PTM:Prolines at the third position of the tripeptide repeating unit (G-X-Y) are hydroxylated in some or all of the chains.,similarity:Belongs to the fibril-associated collagens with interrupted helices (FACIT) family.,subunit:Heterotrimer of an alpha 1(IX), an alpha 2(IX) and an alpha 3(IX) chain.,
Subcellular Location :	Secreted, extracellular space, extracellular matrix .
Expression :	Cartilage,Pancreas,Placenta,Skin,
Sort :	4400
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Immunofluorescence analysis of NIH/3T3 cells, using Collagen IX alpha3 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human tonsil tissue, using Collagen IX alpha3 Antibody. The picture on the right is blocked with the synthesized peptide.