

**COX7a2/3 Polyclonal Antibody**

<b>Catalog No :</b>	YT1079
<b>Reactivity :</b>	Human;Mouse;Rat
<b>Applications :</b>	WB;IHC;IF;ELISA
<b>Target :</b>	COX7a2/3
<b>Gene Name :</b>	COX7A2/COX7A2P2
<b>Protein Name :</b>	Cytochrome c oxidase subunit 7A2/3 mitochondrial
<b>Human Gene Id :</b>	1347
<b>Human Swiss Prot No :</b>	O60397/P14406
<b>Mouse Gene Id :</b>	12866
<b>Rat Gene Id :</b>	29507
<b>Rat Swiss Prot No :</b>	P35171
<b>Immunogen :</b>	The antiserum was produced against synthesized peptide derived from human COX7S/A2. AA range:1-50
<b>Specificity :</b>	COX7a2/3 Polyclonal Antibody detects endogenous levels of COX7a2/3 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000.. IF 1:50-200
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml

---

**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)

---

**Observed Band :** 12kD

---

**Background :** Cytochrome c oxidase, the terminal component of the mitochondrial respiratory chain, catalyzes the electron transfer from reduced cytochrome c to oxygen. This component is a heteromeric complex consisting of three catalytic subunits encoded by mitochondrial genes, and multiple structural subunits encoded by nuclear genes. The mitochondrially-encoded subunits function in electron transfer, while the nuclear-encoded subunits may function in the regulation and assembly of the complex. This nuclear gene encodes polypeptide 2 (liver isoform) of subunit VIIa, with this polypeptide being present in both muscle and non-muscle tissues. In addition to polypeptide 2, subunit VIIa includes polypeptide 1 (muscle isoform), which is present only in muscle tissues, and a related protein, which is present in all tissues. Alternative splicing results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 4 and 14.

---

**Subcellular Location :** Mitochondrion inner membrane .

---

**Tag :** orthogonal

---

**Sort :** 4492

---

**No4 :** 1

---


**Host :** Rabbit

---


**Modifications :** Unmodified

---


## Products Images



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using COX7S/A2 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from rat heart cells, using COX7S/A2 Antibody. The lane on the right is blocked with the synthesized peptide.



The nature compound dehydrocrenatidine exerts potent antihepatocellular carcinoma by destroying mitochondrial complexes in vitro and in vivo 2022 Feb 02. WB Human