

EP-CAM Polyclonal Antibody

Catalog No :	YT1572
Reactivity :	Human;Rat;Mouse;
Applications :	WB;IF;ELISA
Target :	Ep-CAM
Gene Name :	EPCAM
Protein Name :	Epithelial cell adhesion molecule
Human Gene Id :	4072
Human Swiss Prot No :	P16422
Mouse Swiss Prot No :	Q99JW5
Immunogen :	The antiserum was produced against synthesized peptide derived from human TACD1. AA range:116-165
Specificity :	EP-CAM Polyclonal Antibody detects endogenous levels of EP-CAM protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:5000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	34kD

Background : This gene encodes a carcinoma-associated antigen and is a member of a family that includes at least two type I membrane proteins. This antigen is expressed on most normal epithelial cells and gastrointestinal carcinomas and functions as a homotypic calcium-independent cell adhesion molecule. The antigen is being used as a target for immunotherapy treatment of human carcinomas. Mutations in this gene result in congenital tufting enteropathy. [provided by RefSeq, Dec 2008],

Subcellular Location : Lateral cell membrane ; Single-pass type I membrane protein . Cell junction, tight junction . Colocalizes with CLDN7 at the lateral cell membrane and tight junction. .

Expression : Highly and selectively expressed by undifferentiated rather than differentiated embryonic stem cells (ESC). Levels rapidly diminish as soon as ESC's differentiate (at protein levels). Expressed in almost all epithelial cell membranes but not on mesodermal or neural cell membranes. Found on the surface of adenocarcinoma.

Tag : orthogonal


Sort : 5597

No4 : 1


Host : Rabbit

Modifications : Unmodified


Products Images



Western Blot analysis of various cells using EP-CAM Polyclonal Antibody diluted at 1:2000



Western Blot analysis of 3T3 cells using EP-CAM Polyclonal Antibody diluted at 1:2000



Western blot analysis of lysates from Jurkat, HUVEC, and COLO205 cells, using TACD1 Antibody. The lane on the right is blocked with the synthesized peptide.