

GRIP-1 Polyclonal Antibody

Catalog No :	YT2062
Reactivity :	Human;Rat;Mouse;
Applications :	IHC;IF;ELISA
Target :	GRIP-1
Fields :	>>Estrogen signaling pathway;>>Thyroid hormone signaling pathway
Gene Name :	NCOA2
Protein Name :	Nuclear receptor coactivator 2
Human Gene Id :	10499
Human Swiss Prot No :	Q15596
Mouse Swiss Prot No :	Q61026
Immunogen :	The antiserum was produced against synthesized peptide derived from human NCoA2. AA range:702-751
Specificity :	GRIP-1 Polyclonal Antibody detects endogenous levels of GRIP-1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight : 159kD

Background : The protein encoded by this gene functions as a transcriptional coactivator for nuclear hormone receptors, including steroid, thyroid, retinoid, and vitamin D receptors. The encoded protein acts as an intermediary factor for the ligand-dependent activity of these nuclear receptors, which regulate their target genes upon binding of cognate response elements. This gene has been found to be involved in translocations that result in fusions with other genes in various cancers, including the lysine acetyltransferase 6A (KAT6A) gene in acute myeloid leukemia, the ETS variant 6 (ETV6) gene in acute lymphoblastic leukemia, and the hes related family bHLH transcription factor with YRPW motif 1 (HEY1) gene in mesenchymal chondrosarcoma. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016],

Function : disease:Chromosomal aberrations involving NCOA2 may be a cause of acute myeloid leukemias. Inversion inv(8)(p11;q13) generates the MYST3-NCOA2 oncogene, which consists of the N-terminus part of MYST3/MOZ and the C-terminus part of NCOA2/TIF2. MYST3-NCOA2 binds to CREBBP and disrupts its function in transcription activation.,domain:Contains 2 C-terminal transcription activation domains (AD1 and AD2) that can function independently.,domain:Contains four Leu-Xaa-Xaa-Leu-Leu (LXXLL) motifs. The LXXLL motifs are essential for the association with nuclear receptors and are, at least in part, functionally redundant.,domain:The LLXXLXXXL motif is involved in transcriptional coactivation and CREBBP/CBP binding.,function:Transcriptional coactivator for steroid receptors and nuclear receptors. Coactivator of the steroid binding domain (AF-2) but not of the modulating N-terminal domain (AF-1). Requi

Subcellular Location : Nucleus .

Expression : Epithelium,Placenta,Spinal cord,Testis,


Sort : 7116

No4 : 1

Host : Rabbit

Modifications : Unmodified

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4 ° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).