

GlyR β Polyclonal Antibody

Catalog No :	YT1935
Reactivity :	Human;Mouse;Rat
Applications :	WB;ELISA
Target :	GlyR β
Fields :	>>Neuroactive ligand-receptor interaction
Gene Name :	GLRB
Protein Name :	Glycine receptor subunit beta
Human Gene Id :	2743
Human Swiss Prot No :	P48167
Mouse Gene Id :	14658
Mouse Swiss Prot No :	P48168
Rat Gene Id :	25456
Rat Swiss Prot No :	P20781
Immunogen :	The antiserum was produced against synthesized peptide derived from human GLRB. AA range:211-260
Specificity :	GlyR β Polyclonal Antibody detects endogenous levels of GlyR β protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications.

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	56kD
Cell Pathway :	Neuroactive ligand-receptor interaction;
Background :	<p>This gene encodes the beta subunit of the glycine receptor, which is a pentamer composed of alpha and beta subunits. The receptor functions as a neurotransmitter-gated ion channel, which produces hyperpolarization via increased chloride conductance due to the binding of glycine to the receptor. Mutations in this gene cause startle disease, also known as hereditary hyperekplexia or congenital stiff-person syndrome, a disease characterized by muscular rigidity. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Oct 2009],</p>
Function :	<p>disease:Defects in GLRB are a cause of startle disease (STHE) [MIM:149400]; also known as hereditary hyperekplexia or congenital stiff-person syndrome. STHE is a genetically heterogeneous neurologic disorder characterized by muscular rigidity of central nervous system origin, particularly in the neonatal period, and by an exaggerated startle response to unexpected acoustic or tactile stimuli. Inheritance can be autosomal dominant or recessive.,function:The glycine receptor is a neurotransmitter-gated ion channel. Binding of glycine to its receptor increases the chloride conductance and thus produces hyperpolarization (inhibition of neuronal firing).,similarity:Belongs to the ligand-gated ionic channel (TC 1.A.9) family.,subunit:Pentamer composed of alpha and beta subunits. Interacts with GPHN.,</p>
Subcellular Location :	<p>Cell junction, synapse, postsynaptic cell membrane ; Multi-pass membrane protein . Cell junction, synapse . Cell projection, dendrite . Cell membrane ; Multi-pass membrane protein . Cytoplasm . Retained in the cytoplasm upon heterologous expression by itself. Coexpression with GPHN promotes expression at the cell membrane (PubMed:12684523). Coexpression with GLRA1, GLRA2 or GLRA3 promotes expression at the cell membrane. .</p>
Expression :	Brain,Hippocampus,
Sort :	6663
No4 :	1
Host :	Rabbit

Modifications : Unmodified**Products Images**