

## IRAK-2 Polyclonal Antibody

<b>Catalog No :</b>	YT2392
<b>Reactivity :</b>	Human;Rat;Mouse;
<b>Applications :</b>	WB;IHC
<b>Target :</b>	IRAK-2
<b>Fields :</b>	>>Neurotrophin signaling pathway;>>Tuberculosis
<b>Gene Name :</b>	IRAK2
<b>Protein Name :</b>	Interleukin-1 receptor-associated kinase-like 2
<b>Human Gene Id :</b>	3656
<b>Human Swiss Prot No :</b>	O43187
<b>Mouse Swiss Prot No :</b>	Q8CFA1
<b>Immunogen :</b>	Synthesized peptide derived from the Internal region of human IRAK-2.
<b>Specificity :</b>	IRAK-2 Polyclonal Antibody detects endogenous levels of IRAK-2 protein.
<b>Formulation :</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source :</b>	Polyclonal, Rabbit,IgG
<b>Dilution :</b>	WB 1:500-2000;IHC 1:50-300
<b>Purification :</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Concentration :</b>	1 mg/ml
<b>Storage Stability :</b>	-15°C to -25°C/1 year(Do not lower than -25°C)

**Observed Band :** 70kD**Cell Pathway :**Apoptosis\_Inhibition;Apoptosis\_Mitochondrial;Apoptosis\_Overview;Neurotrophin;**Background :**

IRAK2 encodes the interleukin-1 receptor-associated kinase 2, one of two putative serine/threonine kinases that become associated with the interleukin-1 receptor (IL1R) upon stimulation. IRAK2 is reported to participate in the IL1-induced upregulation of NF-kappaB. [provided by RefSeq, Jul 2008],

---

**Function :**


caution:Asn-335 is present instead of the conserved Asp which is expected to be an active site residue. This enzyme has been shown to be catalytically inactive.,domain:The protein kinase domain is predicted to be catalytically inactive.,function:Binds to the IL-1 type I receptor following IL-1 engagement, triggering intracellular signaling cascades leading to transcriptional up-regulation and mRNA stabilization.,similarity:Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family. Pelle subfamily.,similarity:Contains 1 death domain.,similarity:Contains 1 protein kinase domain.,subunit:Interacts with MYD88. IL-1 stimulation leads to the formation of a signaling complex which dissociates from the IL-1 receptor following the binding of PELI1.,tissue specificity:Expressed in spleen, thymus, prostate, lung, liver, skeletal muscle, kidney, pancreas and peripheral blood leuko

---


**Subcellular Location :**nucleus,cytoplasm,cytosol,plasma membrane,endosome membrane,**Expression :**Expressed in spleen, thymus, prostate, lung, liver, skeletal muscle, kidney, pancreas and peripheral blood leukocytes.**Tag :**orthogonal**Sort :**8662**No4 :**1**Host :**Rabbit**Modifications :**Unmodified

---


**Products Images**



Western Blot analysis of various cells using IRAK-2 Polyclonal Antibody diluted at 1:2000



Western Blot analysis of 293 cells using IRAK-2 Polyclonal Antibody diluted at 1:2000



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).