

HBO1 Polyclonal Antibody

Catalog No :	YT2101
Reactivity :	Human;Mouse;Rat
Applications :	WB;IF;ELISA
Target :	HBO1
Gene Name :	KAT7
Protein Name :	Histone acetyltransferase KAT7
Human Gene Id :	11143
Human Swiss Prot No :	O95251
Mouse Gene Id :	217127
Mouse Swiss Prot No :	Q5SVQ0
Rat Gene Id :	303470
Rat Swiss Prot No :	Q810T5
Immunogen :	The antiserum was produced against synthesized peptide derived from human MYST2. AA range:131-180
Specificity :	HBO1 Polyclonal Antibody detects endogenous levels of HBO1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml**Storage Stability :** -15°C to -25°C/1 year(Do not lower than -25°C)**Observed Band :** 75kD**Cell Pathway :** Protein_Acetylation**Background :**

catalytic activity:Acetyl-CoA + histone = CoA + acetylhistone.,domain:The C2HC-type zinc finger is required for interaction with MCM2 and ORC1L.,domain:The N-terminus is involved in transcriptional repression, while the C-terminus mediates AR-interaction.,function:Component of the HBO1 complex which has a histone H4-specific acetyltransferase activity, a reduced activity toward histone H3 and is responsible for the bulk of histone H4 acetylation in vivo. Through chromatin acetylation it may regulate DNA replication and act as a coactivator of TP53-dependent transcription. Specifically represses AR-mediated transcription.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the MYST (SAS/MOZ) family.,similarity:Contains 1 C2HC-type zinc finger.,subunit:Component of the HBO1 complex composed at least of ING4 or ING5, MYTS2/HBO1, EAF6, and one of PHF15, PHF16 and PHF17. Interacts with MCM2 and ORC1L. Interacts with the androgen receptor (AR) in the presence of dihydrotestosterone.,tissue specificity:Ubiquitously expressed, with highest levels in testis.,

Function :

catalytic activity:Acetyl-CoA + histone = CoA + acetylhistone.,domain:The C2HC-type zinc finger is required for interaction with MCM2 and ORC1L.,domain:The N-terminus is involved in transcriptional repression, while the C-terminus mediates AR-interaction.,function:Component of the HBO1 complex which has a histone H4-specific acetyltransferase activity, a reduced activity toward histone H3 and is responsible for the bulk of histone H4 acetylation in vivo. Through chromatin acetylation it may regulate DNA replication and act as a coactivator of TP53-dependent transcription. Specifically represses AR-mediated transcription.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the MYST (SAS/MOZ) family.,similarity:Contains 1 C2HC-type zinc finger.,subunit:Component of the HBO1 complex composed at least of ING4 or ING5, MYTS2/HBO1, EAF6, and one of PHF15, PHF16 an


Subcellular Location :

Nucleus . Chromosome . Chromosome, centromere . Cytoplasm, cytosol . Associates with replication origins specifically during the G1 phase of the cell cycle (PubMed:18832067, PubMed:20129055). Localizes to transcription start sites (PubMed:21753189, PubMed:24065767). Localizes to ultraviolet-induced DNA damage sites following phosphorylation by ATR (PubMed:28719581). Localizes to centromeres in G1 phase (PubMed:27270040). .


Expression : Ubiquitously expressed, with highest levels in testis.**Tag :** hot

Sort :	7254
No4 :	1
Host :	Rabbit
Modifications :	Unmodified


Products Images




Western Blot analysis of various cells using HBO1 Polyclonal Antibody diluted at 1:2000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Western Blot analysis of Jurkat cells using HBO1 Polyclonal Antibody diluted at 1:2000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Immunofluorescence analysis of HUVEC cells, using MYST2 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from Jurkat cells, using MYST2 Antibody. The lane on the right is blocked with the synthesized peptide.