

hnRNP K Polyclonal Antibody

Catalog No :	YT2202
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	hnRNP K
Fields :	>>Spliceosome;>>Viral carcinogenesis;>>MicroRNAs in cancer
Gene Name :	HNRNPK
Protein Name :	Heterogeneous nuclear ribonucleoprotein K
Human Gene Id :	3190
Human Swiss Prot No :	P61978
Mouse Gene Id :	15387
Mouse Swiss Prot No :	P61979
Rat Gene Id :	117282
Rat Swiss Prot No :	P61980
Immunogen :	The antiserum was produced against synthesized peptide derived from human hnRNP K. AA range:250-299
Specificity :	hnRNP K Polyclonal Antibody detects endogenous levels of hnRNP K protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	51kD
Cell Pathway :	Spliceosome;
Background :	<p>This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are RNA binding proteins and they complex with heterogeneous nuclear RNA (hnRNA). These proteins are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and transport. While all of the hnRNPs are present in the nucleus, some seem to shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene is located in the nucleoplasm and has three repeats of KH domains that binds to RNAs. It is distinct among other hnRNP proteins in its binding preference; it binds tenaciously to poly(C). This protein is also thought to have a role during cell cycle progression. Several alternatively spliced transcript variants have</p>
Function :	<p>function:One of the major pre-mRNA-binding proteins. Binds tenaciously to poly(C) sequences. Likely to play a role in the nuclear metabolism of hnRNAs, particularly for pre-mRNAs that contain cytidine-rich sequences. Can also bind poly(C) single-stranded DNA.,mass spectrometry: PubMed:11840567,PTM:Arg-296 and Arg-299 are dimethylated, probably to asymmetric dimethylarginine.,similarity:Contains 1 KH domain.,similarity:Contains 2 KH domains.,similarity:Contains 3 KH domains.,subcellular location:In case of ASFV infection, there is a shift in the localization which becomes predominantly nuclear.,subunit:Interacts with RBM42 and ZIK1 (By similarity). Identified in the spliceosome C complex, at least composed of AQR, ASCC3L1, C19orf29, CDC40, CDC5L, CRNKL1, DDX23, DDX41, DDX48, DDX5, DGCR14, DHX35, DHX38, DHX8, EFTUD2, FRG1, GPATC1, HNRPA1, HNRPA2B1, HNRPA3, HNRPC, HNRPF, HNRPH1, HNRNPK, HNR</p>
Subcellular Location :	Cytoplasm . Nucleus, nucleoplasm . Cell projection, podosome . Recruited to p53/TP53-responsive promoters, in the presence of functional p53/TP53 (PubMed:16360036). In case of ASFV infection, there is a shift in the localization which becomes predominantly nuclear (PubMed:18775702).
Expression :	Brain,Cajal-Retzius cell,Colorectal cancer and surrounding
Tag :	hot


Sort : 7730

No4 : 1


Host : Rabbit

Modifications : Unmodified


Products Images



Western Blot analysis of various cells using hnRNP K Polyclonal Antibody diluted at 1:2000



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.



Western blot analysis of lysates from K562, A549, and HT-29 cells, using hnRNP K Antibody. The lane on the right is blocked with the synthesized peptide.