


LPAAT-θ Polyclonal Antibody


Catalog No :	YT2584
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	LPAAT-θ
Fields :	>>Glycerolipid metabolism;>>Glycerophospholipid metabolism;>>Metabolic pathways
Gene Name :	AGPAT9
Protein Name :	Glycerol-3-phosphate acyltransferase 3
Human Gene Id :	84803
Human Swiss Prot No :	Q53EU6
Mouse Gene Id :	231510
Mouse Swiss Prot No :	Q8C0N2
Rat Gene Id :	305166
Rat Swiss Prot No :	Q4V8J4
Immunogen :	The antiserum was produced against synthesized peptide derived from human AGPAT9. AA range:381-430
Specificity :	LPAAT-θ Polyclonal Antibody detects endogenous levels of LPAAT-θ protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	48kD
Background :	This gene encodes a member of the lysophosphatidic acid acyltransferase protein family. The encoded protein is an enzyme which catalyzes the conversion of glycerol-3-phosphate to lysophosphatidic acid in the synthesis of triacylglycerol. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, Jan 2012],
Function :	catalytic activity:Acyl-CoA + sn-glycerol 3-phosphate = CoA + 1-acyl-sn-glycerol 3-phosphate.,domain:The HXXXXD motif is essential for acyltransferase activity and may constitute the binding site for the phosphate moiety of the glycerol-3-phosphate.,enzyme regulation:Inhibited by N-ethylmaleimide (NEM).,function:Esterifies acyl-group from acyl-ACP to the sn-1 position of glycerol-3-phosphate, an essential step in glycerolipid biosynthesis. Overexpression activates the mTOR pathway.,pathway:Glycerolipid metabolism; triacylglycerol biosynthesis.,pathway:Phospholipid metabolism; CDP-diacylglycerol biosynthesis; CDP-diacylglycerol from sn-glycerol 3-phosphate: step 1/3.,similarity:Belongs to the 1-acyl-sn-glycerol-3-phosphate acyltransferase family.,tissue specificity:Widely expressed. Expressed in liver, kidney, testis, brain, heart, skeletal muscle, thyroid, prostate, thymus and placenta.
Subcellular Location :	Endoplasmic reticulum membrane ; Multi-pass membrane protein .
Expression :	Widely expressed. Expressed in liver, kidney, testis, brain, heart, skeletal muscle, thyroid, prostate, thymus and placenta. Also expressed lung and adipose tissue.
Sort :	9229
No4 :	1
Host :	Rabbit
Modifications :	Unmodified


Products Images




Western Blot analysis of HEPG2 using LPAAT- θ Polyclonal Antibody diluted at 1:1000




Western blot analysis of SH-SY5Y 3T3 lysis using LPAAT- θ antibody. Antibody was diluted at 1:1000



Immunofluorescence analysis of HepG2 cells, using LPAAT- θ Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from Jurkat cells, COLO205 cells, HeLa cells, and HUVEC cells, using PLCH Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).