


IL-21 Polyclonal Antibody

Catalog No :	YT2327
Reactivity :	Human;Mouse
Applications :	WB;ELISA
Target :	IL-21
Fields :	>>Cytokine-cytokine receptor interaction;>>JAK-STAT signaling pathway;>>Th17 cell differentiation;>>Inflammatory bowel disease
Gene Name :	IL21
Protein Name :	Interleukin-21
Human Gene Id :	59067
Human Swiss Prot No :	Q9HBE4
Mouse Gene Id :	60505
Mouse Swiss Prot No :	Q9ES17
Immunogen :	The antiserum was produced against synthesized peptide derived from human IL21. AA range:91-140
Specificity :	IL-21 Polyclonal Antibody detects endogenous levels of IL-21 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:20000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml

Storage Stability :	<u>-15°C to -25°C/1 year(Do not lower than -25°C)</u>
Observed Band :	<u>18kD</u>
Cell Pathway :	<u>Cytokine-cytokine receptor interaction;Jak_STAT;</u>
Background :	<u>This gene encodes a member of the common-gamma chain family of cytokines with immunoregulatory activity. The encoded protein plays a role in both the innate and adaptive immune responses by inducing the differentiation, proliferation and activity of multiple target cells including macrophages, natural killer cells, B cells and cytotoxic T cells. Dysregulation of this gene plays a role in multiple immune-mediated diseases including lupus, psoriasis and chronic inflammatory diseases. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Dec 2011],</u>
Function :	<u>function:Cytokine with immunoregulatory activity. May promote the transition between innate and adaptive immunity. Induces the production of IgG(1) and IgG(3) in B-cells (By similarity). May play a role in proliferation and maturation of natural killer (NK) cells in synergy with IL15. May regulate proliferation of mature B- and T-cells in response to activating stimuli. In synergy with IL15 and IL18 stimulates interferon gamma production in T-cells and NK cells. During T-cell mediated immune response may inhibit dendritic cells (DC) activation and maturation.,similarity:Belongs to the IL-15/IL-21 family.,tissue specificity:Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes.,</u>
Subcellular Location :	<u>Secreted .</u>
Expression :	<u>Expressed in activated CD4-positive T-cells but not in CD8-positive T-cells, B-cells, or monocytes.</u>
Sort :	<u>8480</u>
No4 :	<u>1</u>
Host :	<u>Rabbit</u>
Modifications :	<u>Unmodified</u>

Products Images



Western Blot analysis of various cells using IL-21 Polyclonal Antibody diluted at 1:500