

MAD2L1BP Polyclonal Antibody

Catalog No :	YT2619
Reactivity :	Human;Rat;Mouse;
Applications :	WB;ELISA
Target :	MAD2L1BP
Gene Name :	MAD2L1BP
Protein Name :	MAD2L1-binding protein
Human Gene Id :	9587
Human Swiss Prot No :	Q15013
Mouse Swiss Prot No :	Q9DCX1
Immunogen :	Synthesized peptide derived from MAD2L1BP . at AA range: 10-90
Specificity :	MAD2L1BP Polyclonal Antibody detects endogenous levels of MAD2L1BP protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	31kD

Background : The protein encoded by this gene was identified as a binding protein of the MAD2 mitotic arrest deficient-like 1 (MAD2/MAD2L1). MAD2 is a key component of the spindle checkpoint that delays the onset of anaphase until all the kinetochores are attached to the spindle. This protein may interact with the spindle checkpoint and coordinate cell cycle events in late mitosis. Alternatively spliced transcript variants encoding distinct isoforms have been observed. [provided by RefSeq, Jul 2008],

Function : developmental stage:During the cell cycle, levels increase and then remain constant until late mitosis after which they drop.,function:May function to silence the spindle checkpoint and allow mitosis to proceed through anaphase by binding MAD2L1 after it has become dissociated from the MAD2L1-CDC20 complex.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the MAD2L1BP family.,subcellular location:During early mitosis, unevenly distributed throughout the nucleoplasm. From metaphase to anaphase, concentrated on the spindle.,subunit:Interacts with MAD2L1.,

Subcellular Location : Nucleus. Cytoplasm, cytoskeleton, spindle. During early mitosis, unevenly distributed throughout the nucleoplasm. From metaphase to anaphase, concentrated on the spindle.

Expression : Bone marrow,Brain,Lung,

Sort : 9329

No4 : 1

Host : Rabbit

Modifications : Unmodified

Products Images

