

MAP-9 Polyclonal Antibody

Catalog No: YT2638

Reactivity: Human; Rat; Mouse;

Applications: WB;ELISA

Target: MAP-9

Gene Name: MAP9

Protein Name: Microtubule-associated protein 9

Q49MG5

Q3TRR0

Human Gene Id: 79884

Human Swiss Prot

No:

Mouse Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

MAP9. AA range:121-170

Specificity: MAP-9 Polyclonal Antibody detects endogenous levels of MAP-9 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source : Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 75kD

1/3



Background:

ASAP is a microtubule-associated protein required for spindle function, mitotic progression, and cytokinesis (Saffin et al., 2005 [PubMed 16049101]).[supplied by OMIM, Mar 2008],

Function:

developmental stage:Constitutively expressed during the cell cycle.,function:Involved in organization of the bipolar mitotic spindle. Required for bipolar spindle assembly, mitosis progression and cytokinesis. May act by stabilizing interphase microtubules.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,subcellular location:Localizes to microtubules in interphase, associates with the mitotic spindle during mitosis, localizes to the central body during cytokinesis.,subunit:Binds to purified microtubules via its C-terminus.,

Subcellular Location:


Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, spindle. Localizes to microtubules in interphase, associates with the mitotic spindle during mitosis, localizes to the central body during cytokinesis.

Expression: Amygdala,Bone marrow,Brain,


Sort : 9373

No4:

Products Images



Western blot analysis of MAP9 Antibody. The lane on the right is blocked with the MAP9 peptide.



Western blot analysis of the lysates from HUVECcells using MAP9 antibody.