


MDMX Polyclonal Antibody

Catalog No :	YT2693
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	MDMX
Fields :	>>p53 signaling pathway;>>MicroRNAs in cancer
Gene Name :	MDM4
Protein Name :	Protein Mdm4
Human Gene Id :	4194
Human Swiss Prot No :	O15151
Mouse Gene Id :	17248
Mouse Swiss Prot No :	O35618
Rat Gene Id :	304798
Rat Swiss Prot No :	Q5XIN1
Immunogen :	The antiserum was produced against synthesized peptide derived from human MDM4. AA range:336-385
Specificity :	MDMX Polyclonal Antibody detects endogenous levels of MDMX protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200


Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	45kD
Cell Pathway :	p53;
Background :	<p>This gene encodes a nuclear protein that contains a p53 binding domain at the N-terminus and a RING finger domain at the C-terminus, and shows structural similarity to p53-binding protein MDM2. Both proteins bind the p53 tumor suppressor protein and inhibit its activity, and have been shown to be overexpressed in a variety of human cancers. However, unlike MDM2 which degrades p53, this protein inhibits p53 by binding its transcriptional activation domain. This protein also interacts with MDM2 protein via the RING finger domain, and inhibits the latter's degradation. So this protein can reverse MDM2-targeted degradation of p53, while maintaining suppression of p53 transactivation and apoptotic functions. Alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq, Feb 2011],</p>
Function :	<p>alternative products:Additional isoforms seem to exist, domain:Region I is sufficient for binding p53 and inhibiting its G1 arrest and apoptosis functions. It also binds p73. Region II contains most of a central acidic region and a putative C4-type zinc finger. The RING finger domain which coordinates two molecules of zinc mediates the heterooligomerization with MDM2., function:Inhibits p53- and p73-mediated cell cycle arrest and apoptosis by binding its transcriptional activation domain. Inhibits degradation of MDM2. Can reverse MDM2-targeted degradation of p53 while maintaining suppression of p53 transactivation and apoptotic functions., mass spectrometry: PubMed:11840567, similarity:Belongs to the MDM2/MDM4 family., similarity:Contains 1 RanBP2-type zinc finger., similarity:Contains 1 RING-type zinc finger., similarity:Contains 1 SWIB domain., subunit: Binds to p53, p73 and MDM2., tissue specif</p>
Subcellular Location :	Nucleus.
Expression :	Expressed in all tissues tested with high levels in thymus.
Sort :	9497
No4 :	1
Host :	Rabbit

Modifications : Unmodified


Products Images



Western Blot analysis of 293T cells using MDMX Polyclonal Antibody diluted at 1:1000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using MDM4 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from rat muscle cells, using MDM4 Antibody. The lane on the right is blocked with the synthesized peptide.