


mGluR2 Polyclonal Antibody


Catalog No :	YT2742
Reactivity :	Human;Mouse;Rat
Applications :	IHC;IF;ELISA
Target :	mGluR2
Fields :	>>Phospholipase D signaling pathway;>>Neuroactive ligand-receptor interaction;>>Glutamatergic synapse;>>Cocaine addiction
Gene Name :	GRM2
Protein Name :	Metabotropic glutamate receptor 2
Human Gene Id :	2912
Human Swiss Prot No :	Q14416
Mouse Gene Id :	108068
Mouse Swiss Prot No :	Q14BI2
Rat Gene Id :	24415
Rat Swiss Prot No :	P31421
Immunogen :	The antiserum was produced against synthesized peptide derived from human GRM2. AA range:241-290
Specificity :	mGluR2 Polyclonal Antibody detects endogenous levels of mGluR2 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:5000. Not yet tested in other applications.

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	96kD
Cell Pathway :	Neuroactive ligand-receptor interaction;
Background :	glutamate metabotropic receptor 2(GRM2) Homo sapiens L-glutamate is the major excitatory neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Two transcript variants encoding different isoforms have been found for this gene
Function :	function:Receptor for glutamate. The activity of this receptor is mediated by a G-protein that inhibits adenylate cyclase activity. May mediate suppression of neurotransmission or may be involved in synaptogenesis or synaptic stabilization.,similarity:Belongs to the G-protein coupled receptor 3 family.,subunit:Interacts with GRASP.,tissue specificity:Widely expressed in different regions of the adult brain as well as in fetal brain.,
Subcellular Location :	Cell membrane; Multi-pass membrane protein. Cell junction, synapse . Cell projection, dendrite .
Expression :	Detected in brain cortex (at protein level). Widely expressed in different regions of the adult brain as well as in fetal brain.
Sort :	9614
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Immunofluorescence analysis of LOVO cells, using GRM2 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using GRM2 Antibody. The picture on the right is blocked with the synthesized peptide.