

Laminin α-5 Polyclonal Antibody

| Catalog No : | YT2527 |
|--------------------------|--|
| Reactivity : | Human;Mouse |
| Applications : | WB;IHC;IF;ELISA |
| Target : | LAMA5 |
| Fields : | >>PI3K-Akt signaling pathway;>>Focal adhesion;>>ECM-receptor interaction;>>Toxoplasmosis;>>Amoebiasis;>>Human papillomavirus infection;>>Pathways in cancer;>>Small cell lung cancer |
| Gene Name : | LAMA5 |
| Protein Name : | Laminin subunit alpha-5 |
| Human Gene Id : | 3911 |
| Human Swiss Prot No : | O15230 |
| Mouse Gene Id : | 16776 |
| Mouse Swiss Prot No : | Q61001 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human LAMA5. AA range:2381-2430 |
| Specificity : | Laminin α -5 Polyclonal Antibody detects endogenous levels of Laminin α -5 protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | WB 1:500 - 1:2000. IHC: 1:100-300 ELISA: 1:20000. IF 1:100-300 Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity- |



chromatography using epitope-specific immunogen.

| Concentration : | 1 mg/ml |
|---------------------------|---|
| | |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |
| Observed Band : | 400kD |
| Cell Pathway : | Focal adhesion;ECM-receptor interaction;Pathways in cancer;Small cell lung cancer; |
| Background : | This gene encodes one of the vertebrate laminin alpha chains. Laminins, a family of extracellular matrix glycoproteins, are the major noncollagenous constituent of basement membranes. They have been implicated in a wide variety of biological processes including cell adhesion, differentiation, migration, signaling, neurite outgrowth and metastasis. Laminins are composed of 3 non identical chains: laminin alpha, beta and gamma (formerly A, B1, and B2, respectively) and they form a cruciform structure consisting of 3 short arms, each formed by a different chain, and a long arm composed of all 3 chains. Each laminin chain is a multidomain protein encoded by a distinct gene. The protein encoded by this gene is the alpha-5 subunit of of laminin-10 (laminin-511), laminin-11 (laminin-521) and laminin-15 (laminin-523). [provided by RefSeq, Jun 2013], |
| Function : | domain:Domain G is globular and is part of the major cell-binding site located in the long arm of the laminin heterotrimer.,function:Binding to cells via a high affinity receptor, laminin is thought to mediate the attachment, migration and organization of cells into tissues during embryonic development by interacting with other extracellular matrix components.,similarity:Contains 1 laminin IV type A domain.,similarity:Contains 1 laminin N-terminal domain.,similarity:Contains 22 laminin EGF-like domains.,similarity:Contains 5 laminin G-like domains.,subcellular location:Major component.,subunit:Laminin-15 complex is an heterotrimer composed of three chains (alpha-5/beta-2/gamma-3) which are bound to each other by disulfide bonds into a cross-shaped molecule comprising one long and three short arms with globules at each end.,tissue specificity:Expressed in heart, lung, kidney, skeletal mus |
| Subcellular Location : | Secreted, extracellular space, extracellular matrix, basement membrane. Major component. |
| Expression : | Expressed in heart, lung, kidney, skeletal muscle, pancreas, retina and placenta. Little or no expression in brain and liver. |
| Tag : | hot |
| Sort : | 1356 |



| Best Tools for immunology Research | | |
|------------------------------------|------------|--|
| No3 : | ab14509 | |
| No4 : | 1 | |
| Host : | Rabbit | |
| Modifications : | Unmodified | |

Products Images



LAMA5

Immunofluorescence analysis of A549. 1, primary Antibody was diluted at 1:200(4°C overnight). 2, Goat Anti Rabbit IgG (H&L) - Alexa Fluor 488 Secondary antibody was diluted at 1:1000(room temperature, 50min).3, Picture B: DAPI(blue) 10min.

Western Blot analysis of RAW HepG2 A549 cells using Laminin α -5 Polyclonal Antibody




25---


70-55-40-35-

15----





Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using LAMA5 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night