

MASTL Polyclonal Antibody

YT2661 Catalog No:

Reactivity: Human; Mouse

Applications: WB;ELISA

Target: THC2

Gene Name: MASTL

Protein Name: Serine/threonine-protein kinase greatwall

Q96GX5

Q8C0P0

Human Gene Id: 84930

Human Swiss Prot

No:

Mouse Swiss Prot

Immunogen:

No:

Synthesized peptide derived from the C-terminal region of human MASTL.

Specificity: MASTL Polyclonal Antibody detects endogenous levels of MASTL protein.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Polyclonal, Rabbit, IgG Source:

Dilution: WB 1:500 - 1:2000. ELISA: 1:40000. Not yet tested in other applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

-15°C to -25°C/1 year(Do not lower than -25°C) **Storage Stability:**

Observed Band: 100kD



Background:

This gene encodes a microtubule-associated serine/threonine kinase. Mutations at this locus have been associated with autosomal dominant thrombocytopenia, also known as thrombocytopenia-2. Alternatively spliced transcript variants have been described for this locus. [provided by RefSeq, Feb 2010],

Function:

catalytic activity:ATP + a protein = ADP + a phosphoprotein.,disease:Defects in MASTL are the cause of thrombocytopenia type 2 (THC2) [MIM:188000]. Thrombocytopenia is defined by a decrease in the number of platelets in circulating blood, resulting in the potential for increased bleeding and decreased ability for clotting.,function:Putative serine/threonine kinase which may be involved in megakaryocyte differentiation.,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family.,similarity:Contains 1 AGC-kinase C-terminal domain.,similarity:Contains 1 protein kinase domain.,

Subcellular Location:

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Nucleus. Cleavage furrow. During interphase is mainly nuclear, upon nuclear envelope breakdown localizes at the cytoplasm and during mitosis at the centrosomes. Upon mitotic exit moves to the cleavage furrow.

Expression: Epithelium, Placenta,

Sort : 9417

No4:

Host: Rabbit

Modifications: Unmodified

Products Images

2/2