


mGluR-4 Polyclonal Antibody


Catalog No :	YT2744
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	mGluR-4
Fields :	>>Phospholipase D signaling pathway;>>Neuroactive ligand-receptor interaction;>>Glutamatergic synapse;>>Taste transduction
Gene Name :	GRM4
Protein Name :	Metabotropic glutamate receptor 4
Human Gene Id :	2914
Human Swiss Prot No :	Q14833
Mouse Gene Id :	268934
Mouse Swiss Prot No :	Q68EF4
Rat Gene Id :	24417
Rat Swiss Prot No :	P31423
Immunogen :	The antiserum was produced against synthesized peptide derived from human mGluR4. AA range:851-900
Specificity :	mGluR-4 Polyclonal Antibody detects endogenous levels of mGluR-4 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000.. IF 1:50-200

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	100kD
Cell Pathway :	Neuroactive ligand-receptor interaction;Taste transduction;
Background :	glutamate metabotropic receptor 4(GRM4) Homo sapiens L-glutamate is the major excitatory neurotransmitter in the central nervous system and activates both ionotropic and metabotropic glutamate receptors. Glutamatergic neurotransmission is involved in most aspects of normal brain function and can be perturbed in many neuropathologic conditions. The metabotropic glutamate receptors are a family of G protein-coupled receptors, that have been divided into 3 groups on the basis of sequence homology, putative signal transduction mechanisms, and pharmacologic properties. Group I includes GRM1 and GRM5 and these receptors have been shown to activate phospholipase C. Group II includes GRM2 and GRM3 while Group III includes GRM4, GRM6, GRM7 and GRM8. Group II and III receptors are linked to the inhibition of the cyclic AMP cascade but differ in their agonist selectivities. Several transcript variants encoding different isoforms have been found for this
Function :	function:Receptor for glutamate. The activity of this receptor is mediated by a G-protein that inhibits adenylate cyclase activity.,similarity:Belongs to the G-protein coupled receptor 3 family.,subunit:Interacts with PICK1.,tissue specificity:Strongly expressed in the cerebellum. Expressed at low levels in hippocampus, hypothalamus and thalamus. No expression detected in liver.,
Subcellular Location :	Cell membrane ; Multi-pass membrane protein .
Expression :	Strongly expressed in the cerebellum. Expressed at low levels in hippocampus, hypothalamus and thalamus. No expression detected in liver.
Tag :	orthogonal
Sort :	9616
No4 :	1
Host :	Rabbit
Modifications :	Unmodified


Products Images




Western Blot analysis of various cells using mGluR-4 Polyclonal Antibody diluted at 1:500



Western Blot analysis of 293 cells using mGluR-4 Polyclonal Antibody diluted at 1:500



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using mGluR4 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from 293 cells, treated with Forskolin 40nM 30', using mGluR4 Antibody. The lane on the right is blocked with the synthesized peptide.