


Olfactory receptor 10K1/2 Polyclonal Antibody


Catalog No :	YT3262
Reactivity :	Human;Rat;Mouse;
Applications :	WB;IF;ELISA
Target :	Olfactory receptor 10K1/2
Fields :	>>Olfactory transduction
Gene Name :	OR10K1/OR10K2
Protein Name :	Olfactory receptor 10K1/2
Human Gene Id :	391109/391107
Human Swiss Prot No :	Q8NGX5/Q6IF99
Immunogen :	The antiserum was produced against synthesized peptide derived from human OR10K1/10K2. AA range:56-105
Specificity :	Olfactory receptor 10K1/2 Polyclonal Antibody detects endogenous levels of Olfactory receptor 10K1/2 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band :	35kD
Cell Pathway :	Olfactory transduction;
Background :	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],
Function :	function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
Subcellular Location :	Cell membrane; Multi-pass membrane protein.
Sort :	11092
No4 :	1
Host :	Rabbit
Modifications :	Unmodified


Products Images



Immunofluorescence analysis of LOVO cells, using OR10K1/10K2 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HepG2 cells, using OR10K1/10K2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HUVEC cells using OR10K1/10K2 antibody.