

Olfactory receptor 2B8 Polyclonal Antibody

Catalog No :	YT3300
Reactivity :	Human;Rat;Mouse;
Applications :	WB;IF;ELISA
Target :	Olfactory receptor 2B8
Gene Name :	OR2B8P
Protein Name :	Putative olfactory receptor 2B8
Human Gene Id :	65944
Human Swiss Prot No :	P59922
Immunogen :	The antiserum was produced against synthesized peptide derived from human OR2B8. AA range:61-110
Specificity :	Olfactory receptor 2B8 Polyclonal Antibody detects endogenous levels of Olfactory receptor 2B8 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	38kD

Background : Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Function : caution:Could be the product of a pseudogene.,function:Odorant receptor
.,similarity:Belongs to the G-protein coupled receptor 1 family.,

Subcellular Location : Cell membrane; Multi-pass membrane protein.


Sort : 11130


No4 : 1

Host : Rabbit

Modifications : Unmodified

Products Images





Western blot analysis of the lysates from HT-29 cells using OR2B8 antibody.