


## Neuro D Polyclonal Antibody


|                              |   |
|------------------------------|---|
| <b>Catalog No :</b>          | YT3061  |
| <b>Reactivity :</b>          | Human;Mouse;Rat   |
| <b>Applications :</b>        | WB;ELISA;IHC  |
| <b>Target :</b>              | Neuro D   |
| <b>Fields :</b>              | >>Maturity onset diabetes of the young  |
| <b>Gene Name :</b>           | NEUROD1   |
| <b>Protein Name :</b>        | Neurogenic differentiation factor 1   |
| <b>Human Gene Id :</b>       | 4760  |
| <b>Human Swiss Prot No :</b> | Q13562  |
| <b>Mouse Gene Id :</b>       | 18012   |
| <b>Mouse Swiss Prot No :</b> | Q60867  |
| <b>Rat Gene Id :</b>         | 29458   |
| <b>Rat Swiss Prot No :</b>   | Q64289  |
| <b>Immunogen :</b>           | The antiserum was produced against synthesized peptide derived from human NEUROD1. AA range:240-289 |
| <b>Specificity :</b>         | Neuro D Polyclonal Antibody detects endogenous levels of Neuro D protein.                           |
| <b>Formulation :</b>         | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.                             |
| <b>Source :</b>              | Polyclonal, Rabbit,IgG  |
| <b>Dilution :</b>            | WB 1:500-2000;IHC 1:50-300; ELISA 2000-20000  |

|                               |   |
|-------------------------------|---|
| <b>Purification :</b>         | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |
| <b>Concentration :</b>        | 1 mg/ml   |
| <b>Storage Stability :</b>    | -15°C to -25°C/1 year(Do not lower than -25°C)  |
| <b>Observed Band :</b>        | 36kD  |
| <b>Cell Pathway :</b>         | Maturity onset diabetes of the young;   |
| <b>Background :</b>           | This gene encodes a member of the NeuroD family of basic helix-loop-helix (bHLH) transcription factors. The protein forms heterodimers with other bHLH proteins and activates transcription of genes that contain a specific DNA sequence known as the E-box. It regulates expression of the insulin gene, and mutations in this gene result in type II diabetes mellitus. [provided by RefSeq, Jul 2008],  |
| <b>Function :</b>             | disease:Defects in NEUROD1 are the cause of maturity onset diabetes of the young type 6 (MODY6) [MIM:606394]. MODY [MIM:606391] is characterized by an autosomal dominant mode of inheritance, onset during young adulthood and a primary defect in insulin secretion.,function:Differentiation factor required for dendrite morphogenesis and maintenance in the cerebellar cortex. Transcriptional activator. Binds to the insulin gene E-box.,PTM:Phosphorylated. In islet cells, phosphorylated on Ser-274 upon glucose stimulation; which may be required for nuclear localization. In activated neurons, phosphorylated on Ser-335; which promotes dendritic growth.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,subunit:Efficient DNA binding requires dimerization with another bHLH protein. Heterodimer with TCF3/E47. Interacts with RREB1., |
| <b>Subcellular Location :</b> | Cytoplasm . Nucleus . In pancreatic islet cells, shuttles to the nucleus in response to glucose stimulation (By similarity). Colocalizes with NR0B2 in the nucleus. .   |
| <b>Expression :</b>           | Eye,Retina,Rhabdomyosarcoma,  |
| <b>Sort :</b>                 | 10711   |
| <b>No4 :</b>                  | 1   |
| <b>Host :</b>                 | Rabbit  |
| <b>Modifications :</b>        | Unmodified  |

## Products Images



Western blot analysis of lysates from HeLa cells, using Neuro D Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).