

Nidogen Polyclonal Antibody

Catalog No :	YT3126		
Reactivity :	Human;Mouse;Rat		
Applications :	WB;ELISA		
Target :	Nidogen		
Gene Name :	NID1		
Protein Name :	Nidogen-1		
Human Gene Id :	4811		
Human Swiss Prot	P14543		
No : Mouse Gene Id :	18073		
Mouse Swiss Prot	P10493		
No : Rat Swiss Prot No :	P08460		
Immunogen :	The antiserum was produced against synthesized peptide derived from human NID1. AA range:219-268		
Specificity :	Nidogen Polyclonal Antibody detects endogenous levels of Nidogen protein.		
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.		
Source :	Polyclonal, Rabbit,IgG		
Dilution :	WB 1:500 - 1:2000. ELISA: 1:20000. Not yet tested in other applications.		
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.		
Concentration :	1 mg/ml		



Dest fools for infinitutiology Research			
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)		
Observed Band :	136kD		
Background :	nidogen 1(NID1) Homo sapiens This gene encodes a member of the nidogen family of basement membrane glycoproteins. The protein interacts with several other components of basement membranes, and may play a role in cell interactions with the extracellular matrix. [provided by RefSeq, Jul 2008],		
Function :	function:Sulfated glycoprotein widely distributed in basement membranes and tightly associated with laminin. Also binds to collagen IV and perlecan. It probably has a role in cell-extracellular matrix interactions.,PTM:N- and O- glycosylated.,similarity:Contains 1 NIDO domain.,similarity:Contains 1 nidogen G2 beta-barrel domain.,similarity:Contains 1 thyroglobulin type-1 domain.,similarity:Contains 4 LDL-receptor class B repeats.,similarity:Contains 6 EGF-like domains.,subunit:Interacts with FBLN1 and LGALS3BP.,		
Subcellular Location :	Secreted, extracellular space, extracellular matrix, basement membrane.		
Expression :	Brain, Placenta, Testis,		
Sort :	10850		
No4 :	1		
Host :	Rabbit		
Modifications :	Unmodified		

Products Images				
к562 к562 170		Western blot analysis of lysates from K562 cells, using NID1 Antibody. The lane on the right is blocked with the synthesized peptide.		
NID1	130			
	95			
	72 (kD)			