

p14ARF Polyclonal Antibody

Catalog No :	YT3490
Reactivity :	Human;Rat;Mouse;
Applications :	WB;IHC;IF;ELISA
Target :	p14ARF
Gene Name :	CDKN2A
Protein Name :	Cyclin-dependent kinase inhibitor 2A isoform 4
Human Gene Id :	1029
Human Swiss Prot No :	Q8N726
Mouse Swiss Prot No :	Q64364
Immunogen :	The antiserum was produced against synthesized peptide derived from human p14 ARF. AA range:71-120
Specificity :	p14 Polyclonal Antibody detects endogenous levels of p14 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	18kD

Cell Pathway : Stem cell pathway; Cell Cycle

Background : CDKN2A generates several transcript variants which differ in their first exons. At least three alternatively spliced variants encoding distinct proteins have been reported, two of which encode structurally related isoforms known to function as inhibitors of CDK4 kinase. The remaining transcript includes an alternate first exon located 20 Kb upstream of the remainder of the gene; this transcript contains an alternate open reading frame (ARF) that specifies a protein which is structurally unrelated to the products of the other variants. This ARF product functions as a stabilizer of the tumor suppressor protein p53 as it can interact with, and sequester, the E3 ubiquitin-protein ligase MDM2, a protein responsible for the degradation of p53. In spite of the structural and functional differences, the CDK inhibitor isoforms and the ARF product encoded by CDKN2A, through the regulatory roles of CDK4 and p53 in cell cycle G1 progression, share a common functionality in cell cycle G1 control. CDKN2A is frequently mutated or deleted in a wide variety of tumors, and is known to be an important tumor suppressor gene.

Subcellular Location : Nucleus, nucleolus . Nucleus, nucleoplasm .; [Isoform smARF]: Mitochondrion .

Sort : 11362

No4 : 1


Host : Rabbit

Modifications : Unmodified


Products Images




Western Blot analysis of various cells using p14 Polyclonal Antibody diluted at 1:500



Immunofluorescence analysis of HeLa cells, using p14 ARF Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human placenta tissue, using p14 ARF Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa cells, using p14 ARF Antibody. The lane on the right is blocked with the synthesized peptide.