

PARP-4 Polyclonal Antibody

Catalog No :	YT3596
Reactivity :	Human;Rat;Mouse;
Applications :	IHC;IF;ELISA
Target :	PARP-4
Fields :	>>Base excision repair;>>Apoptosis
Gene Name :	PARP4
Protein Name :	Poly [ADP-ribose] polymerase 4
Human Gene Id :	143
Human Swiss Prot No :	Q9UKK3
Immunogen :	The antiserum was produced against synthesized peptide derived from human PARP4. AA range:1151-1200
Specificity :	PARP-4 Polyclonal Antibody detects endogenous levels of PARP-4 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	IHC 1:100 - 1:300. ELISA: 1:10000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	193kD

Cell Pathway : Base excision repair;

Background : This gene encodes poly(ADP-ribosyl)transferase-like 1 protein, which is capable of catalyzing a poly(ADP-ribosyl)ation reaction. This protein has a catalytic domain which is homologous to that of poly (ADP-ribosyl) transferase, but lacks an N-terminal DNA binding domain which activates the C-terminal catalytic domain of poly (ADP-ribosyl) transferase. Since this protein is not capable of binding DNA directly, its transferase activity may be activated by other factors such as protein-protein interaction mediated by the extensive carboxyl terminus. [provided by RefSeq, Jul 2008],

Function : catalytic activity:NAD(+) + (ADP-D-ribosyl)(n)-acceptor = nicotinamide + (ADP-D-ribosyl)(n+1)-acceptor.,similarity:Contains 1 BRCT domain.,similarity:Contains 1 PARP alpha-helical domain.,similarity:Contains 1 PARP catalytic domain.,similarity:Contains 1 VWFA domain.,subcellular location:Also found in the nucleus, associated with mitotic spindles.,subunit:Component of the vault ribonucleoprotein particle, at least composed of MVP, PARP4 and one or more vault RNAs (vRNAs). Binds to MVP. Associates with TEP1.,tissue specificity:Widely expressed; the highest levels are in the kidney; also detected in heart, placenta, lung, liver, skeletal muscle, spleen, leukocytes and pancreas.,

Subcellular Location : Cytoplasm . Nucleus . Cytoplasm, cytoskeleton, spindle . Also found in the nucleus, associated with mitotic spindles. .

Expression : Widely expressed; the highest levels are in the kidney; also detected in heart, placenta, lung, liver, skeletal muscle, spleen, leukocytes and pancreas.


Sort : 11642

No4 : 1


Host : Rabbit

Modifications : Unmodified

Products Images



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using PARP4 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HT-29 cells using FAF1 antibody.