

Olfactory receptor 5K3 Polyclonal Antibody

Catalog No: YT3409

Reactivity: Human

Applications: WB;IF;ELISA

Target: Olfactory receptor 5K3

Fields: >>Olfactory transduction

Gene Name: OR5K3

Protein Name: Olfactory receptor 5K3

A6NET4

Human Gene Id: 403277

Human Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

OR5K3. AA range:232-281

Specificity: Olfactory receptor 5K3 Polyclonal Antibody detects endogenous levels of

Olfactory receptor 5K3 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:5000. Not yet tested in other

applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/3



Observed Band: 37kD

Background : olfactory receptor family 5 subfamily K member 3(OR5K3) Homo sapiens

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR)

arising from single coding-exon genes. Olfactory receptors share a

7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by

RefSeq, Jul 2008],

Function: function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor

Cell membrane; Multi-pass membrane protein.

1 family.,

Subcellular


Location:


Sort: 11238

No4:

Host: Rabbit

Modifications: Unmodified





Western blot analysis of the lysates from HepG2 cells using OR5K3 antibody.