

Olfactory receptor 5M11 Polyclonal Antibody

Catalog No: YT3413

Reactivity: Human

Applications: WB;IF;ELISA

Target: Olfactory receptor 5M11

Fields: >>Olfactory transduction

Gene Name: OR5M11

Protein Name: Olfactory receptor 5M11

Human Gene Id: 219487

Human Swiss Prot

No:

s Prot Q96RB7

Immunogen: The antiserum was produced against synthesized peptide derived from human

OR5M11. AA range:233-282

Specificity: Olfactory receptor 5M11 Polyclonal Antibody detects endogenous levels of

Olfactory receptor 5M11 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:5000. Not yet tested in other

applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/3



Observed Band: 34kD

Cell Pathway: Olfactory transduction;

Background: Olfactory receptors interact with odorant molecules in the nose, to initiate a

neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR)

arising from single coding-exon genes. Olfactory receptors share a

7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated

transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by

RefSeq, Jul 2008],

Function: function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor

Cell membrane; Multi-pass membrane protein.

1 family.,

Subcellular

Location:


Sort: 11242

No4:

Host: Rabbit


Modifications: Unmodified

Products Images




Western Blot analysis of various cells using Olfactory receptor 5M11 Polyclonal Antibody





Western blot analysis of lysates from HUVEC cells, using OR5M11 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from K562 cells using OR5M11 antibody.