

Olfactory receptor 6C2 Polyclonal Antibody

Catalog No: YT3424

Reactivity: Human

Applications: WB;IF;ELISA

Target: Olfactory receptor 6C2

Fields: >>Olfactory transduction

Gene Name: OR6C2

Protein Name : Olfactory receptor 6C2

Q9NZP2

Human Gene Id: 341416

Human Swiss Prot

No:

<u>____</u>

Immunogen: The antiserum was produced against synthesized peptide derived from human

OR6C2. AA range:231-280

Specificity: Olfactory receptor 6C2 Polyclonal Antibody detects endogenous levels of

Olfactory receptor 6C2 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other

applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/3



Observed Band: 36kD

Cell Pathway : Olfactory transduction;

Background: olfactory receptor family 6 subfamily C member 2(OR6C2) Homo sapiens

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR)

arising from single coding-exon genes. Olfactory receptors share a

7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated

transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by

RefSeq, Jul 2008],

Function: function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor

Cell membrane; Multi-pass membrane protein.

1 family.,

11253

Subcellular Location:

Sort:


ocation.

No4:


Host: Rabbit

Modifications: Unmodified

Products Images



Western blot analysis of lysates from HT-29 cells, using OR6C2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using OR6C2 antibody.