

Olfactory receptor 6C70 Polyclonal Antibody

Catalog No: YT3427

Reactivity: Human

Applications: WB;ELISA

Target: Olfactory receptor 6C70

Fields: >>Olfactory transduction

Gene Name: OR6C70

Protein Name: Olfactory receptor 6C70

A6NIJ9

Human Gene Id: 390327

Human Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

OR6C70. AA range:231-280

Specificity: Olfactory receptor 6C70 Polyclonal Antibody detects endogenous levels of

Olfactory receptor 6C70 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. ELISA: 1:5000. Not yet tested in other applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 35kD

1/3



Cell Pathway: Olfactory transduction;

Background : Olfactory receptors interact with odorant molecules in the nose, to initiate a

neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR)

arising from single coding-exon genes. Olfactory receptors share a

7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by

RefSeq, Jul 2008],

Function: function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor

1 family.,

Subcellular Location:

Cell membrane; Multi-pass membrane protein.

Tag: hot


Sort : 11256

No4:


Host: Rabbit

Modifications: Unmodified

Products Images



Western blot analysis of lysates from Jurkat cells, using OR6C70 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HeLa cells using OR6C70 antibody.