

Olfactory receptor 6J1 Polyclonal Antibody

Catalog No: YT3428

Reactivity: Human

Applications: WB;IF;ELISA

Target: Olfactory receptor 6J1

Fields: >>Olfactory transduction

Gene Name: OR6J1

Protein Name: Olfactory receptor 6J1

Q8NGC5

Human Gene Id: 79549

Human Swiss Prot

No:

Immunogen: The antiserum was produced against synthesized peptide derived from human

OR6J1. AA range:231-280

Specificity: Olfactory receptor 6J1 Polyclonal Antibody detects endogenous levels of

Olfactory receptor 6J1 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution : WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other

applications.

Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

1/3

Observed Band: 39

39kD

Background:

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Function:

function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor

1 family.,

Subcellular Location:

Cell membrane; Multi-pass membrane protein.


Sort : 11257

No4:


Host: Rabbit

Modifications: Unmodified


Products Images



Immunofluorescence analysis of MCF7 cells, using OR6J1 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HUVEC and MCF-7 cells, using OR6J1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using OR6J1 antibody.