

Olfactory receptor 8B2/3 Polyclonal Antibody

Catalog No :	YT3439
Reactivity :	Human
Applications :	WB;IF;ELISA
Target :	Olfactory receptor 8B2/3
Fields :	>>Olfactory transduction
Gene Name :	OR8B2/OR8B3
Protein Name :	Olfactory receptor 8B2/3
Human Gene Id :	26595
Human Swiss Prot No :	Q96RD0/Q8NKG8
Immunogen :	The antiserum was produced against synthesized peptide derived from human OR8B2/8B3. AA range:199-248
Specificity :	Olfactory receptor 8B2/3 Polyclonal Antibody detects endogenous levels of Olfactory receptor 8B2/3 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 35kD

Background : olfactory receptor family 8 subfamily B member 2(OR8B2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],

Function : function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,

Subcellular Location : Cell membrane; Multi-pass membrane protein.


Sort : 11268

No4 : 1


Host : Rabbit

Modifications : Unmodified

Products Images



Western blot analysis of lysates from Jurkat and HT-29 cells, using OR8B2/8B3 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from K562 cells using OR8B2/B3 antibody.