


ORAOV1 Polyclonal Antibody

| Catalog No : | YT3468 |
|----------------------------------|---|
| Reactivity : | Human;Mouse |
| Applications : | WB;IHC;IF;ELISA |
| Target : | ORAOV1 |
| Gene Name : | ORAOV1 |
| Protein Name : | Oral cancer-overexpressed protein 1 |
| Human Gene Id : | 220064 |
| Human Swiss Prot | Q8WV07 |
| No : Mouse Swiss Prot No : | Q8CH62 |
| Immunogen : | The antiserum was produced against synthesized peptide derived from human ORAV1. AA range:81-130 |
| Specificity : | ORAOV1 Polyclonal Antibody detects endogenous levels of ORAOV1 protein. |
| Formulation : | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source : | Polyclonal, Rabbit,IgG |
| Dilution : | WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other applications. |
| Purification : | The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen. |
| Concentration : | 1 mg/ml |
| Storage Stability : | -15°C to -25°C/1 year(Do not lower than -25°C) |
| Observed Band : | 15kD |




| • | |
|---------------------------|--|
| Background : | miscellaneous:Amplified and overexpressed in oral cancer cells.,similarity:Belongs to the ORAOV1 family.,tissue specificity:Widely expressed. Highly expressed in placenta, kidney and skeletal muscle., |
| Function : | miscellaneous:Amplified and overexpressed in oral cancer cells.,similarity:Belongs to the ORAOV1 family.,tissue specificity:Widely expressed. Highly expressed in placenta, kidney and skeletal muscle., |
| Subcellular Location : | Nucleus . |
| Expression : | Widely expressed. Highly expressed in placenta, kidney and skeletal muscle. |
| Sort : | 11314 |
| No4 : | 1 |
| Host : | Rabbit |
| Modifications : | Unmodified |



Products Images

Western Blot analysis of various cells using ORAOV1 Polyclonal Antibody diluted at 1:2000





Western blot analysis of lysates from 293 and HeLa cells, using ORAV1 Antibody. The lane on the right is blocked with the synthesized peptide.

Immunohistochemical analysis of paraffin-embedded human spleen. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).