

Protocadherin-11 Polyclonal Antibody

Catalog No :	YT3863
Reactivity :	Human;Rat;Mouse;
Applications :	IHC;IF;ELISA
Target :	Protocadherin-11
Gene Name :	PCDH11X/PCDH11Y
Protein Name :	Protocadherin-11 X/Y-linked
Human Gene Id :	83259/27328
Human Swiss Prot No :	Q9BZA8/Q9BZA7
Immunogen :	The antiserum was produced against synthesized peptide derived from human PCDH-X/Y. AA range:531-580
Specificity :	Protocadherin-11 Polyclonal Antibody detects endogenous levels of Protocadherin-11 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Molecularweight :	147kD

Background : This gene belongs to the protocadherin family, a subfamily of the cadherin superfamily. The encoded protein consists of an extracellular domain containing seven cadherin repeats, a transmembrane domain, and a cytoplasmic tail that differs from those of the classical cadherins. This gene is located on the Y chromosome in a block of X/Y homology and is very closely related to its paralog on the X chromosome. The protein is thought to play a role in cell-cell recognition during development of the central nervous system. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2013],

Function : alternative products:Additional isoforms seem to exist,disease:A chromosomal aberration involving PCDH11Y is a cause of multiple congenital abnormalities, including severe bilateral vesicoureteral reflux (VUR) with ureterovesical junction defects. Translocation t(Y;3)(p11;p12) with ROBO2.,function:Potential calcium-dependent cell-adhesion protein.,similarity:Contains 7 cadherin domains.,subunit:Interacts with CTNNB1.,tissue specificity:Expressed strongly in fetal brain and brain (cortex, amygdala, thalamus, substantia nigra, hippocampus, caudate nucleus and corpus callosum). Expressed at low level in testis. Expressed in apoptosis-resistant cells.,

Subcellular Location : Cell membrane ; Single-pass type I membrane protein .

Expression : Expressed strongly in fetal brain and brain (cortex, amygdala, thalamus, substantia nigra, hippocampus, caudate nucleus and corpus callosum). Expressed at low level in testis. Expressed in apoptosis-resistant cells.


Sort : 13074

No4 : 1


Host : Rabbit

Modifications : Unmodified

Products Images



Immunofluorescence analysis of HepG2 cells, using PCDH-X/Y Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using PCDH-X/Y Antibody. The picture on the right is blocked with the synthesized peptide.