

PHLDA3 Polyclonal Antibody

Catalog No :	YT3703
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	PHLDA3
Gene Name :	PHLDA3
Protein Name :	Pleckstrin homology-like domain family A member 3
Human Gene Id :	23612
Human Swiss Prot No :	Q9Y5J5
Mouse Gene Id :	27280
Mouse Swiss Prot No :	Q9WV95
Rat Gene Id :	363989
Rat Swiss Prot No :	Q5PQT7
Immunogen :	The antiserum was produced against synthesized peptide derived from human PHLDA3. AA range:21-70
Specificity :	PHLDA3 Polyclonal Antibody detects endogenous levels of PHLDA3 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.

Concentration : 1 mg/ml

Storage Stability : -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band : 15kD

Background : domain:The PH domain binds phosphoinositides with a broad specificity. It competes with the PH domain of AKT1 and directly interferes with AKT1 binding to phosphatidylinositol 4,5-bisphosphate (PIP2) and phosphatidylinositol 3,4,5-triphosphate (PIP3), preventing AKT1 association to membrane lipids and subsequent activation of AKT1 signaling.,function:p53/TP53-regulated repressor of Akt/AKT1 signaling. Represses AKT1 by preventing AKT1-binding to membrane lipids, thereby inhibiting AKT1 translocation to the cellular membrane and activation. Contributes to p53/TP53-dependent apoptosis by repressing AKT1 activity. Its directs transcription regulation by p53/TP53 may explain how p53/TP53 can negatively regulate AKT1. May acts as a tumor suppressor.,induction:By p53/TP53; expression is directly activated by p53/TP53. p53/TP53 phosphorylation on 'Ser-15' is required to activate the PHLDA3 promoter.,miscellaneous:PHLDA3 genomic locus is frequently observed in primary lung cancers, suggesting a role in tumor suppression.,similarity:Belongs to the PHLDA3 family.,similarity:Contains 1 PH domain.,tissue specificity:Widely expressed with lowest expression in liver and spleen.,

Function : domain:The PH domain binds phosphoinositides with a broad specificity. It competes with the PH domain of AKT1 and directly interferes with AKT1 binding to phosphatidylinositol 4,5-bisphosphate (PIP2) and phosphatidylinositol 3,4,5-triphosphate (PIP3), preventing AKT1 association to membrane lipids and subsequent activation of AKT1 signaling.,function:p53/TP53-regulated repressor of Akt/AKT1 signaling. Represses AKT1 by preventing AKT1-binding to membrane lipids, thereby inhibiting AKT1 translocation to the cellular membrane and activation. Contributes to p53/TP53-dependent apoptosis by repressing AKT1 activity. Its directs transcription regulation by p53/TP53 may explain how p53/TP53 can negatively regulate AKT1. May acts as a tumor suppressor.,induction:By p53/TP53; expression is directly activated by p53/TP53. p53/TP53 phosphorylation on 'Ser-15' is required to activate the PHLDA3 prom

Subcellular Location : Cytoplasm . Membrane ; Peripheral membrane protein .

Expression : Widely expressed with lowest expression in liver and spleen.

Sort : 11895

No4 : 1

Host : Rabbit


Unmodified

Modifications :

Products Images



Western Blot analysis of various cells using PHLDA3 Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using PHLDA3 Antibody. The picture on the right is blocked with the synthesized peptide.