


Rad23A Polyclonal Antibody


Catalog No :	YT3961
Reactivity :	Human;Mouse
Applications :	WB;IHC
Target :	Rad23A
Fields :	>>Nucleotide excision repair;>>Protein processing in endoplasmic reticulum
Gene Name :	RAD23A
Protein Name :	UV excision repair protein RAD23 homolog A
Human Gene Id :	5886
Human Swiss Prot No :	P54725
Mouse Swiss Prot No :	P54726
Immunogen :	The antiserum was produced against synthesized peptide derived from human RAD23A. AA range:91-140
Specificity :	Rad23A Polyclonal Antibody detects endogenous levels of Rad23A protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000;IHC 1:50-300
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band :	48kD
Cell Pathway :	Nucleotide excision repair;
Background :	<p>The protein encoded by this gene is one of two human homologs of <i>Saccharomyces cerevisiae</i> Rad23, a protein involved in nucleotide excision repair. Proteins in this family have a modular domain structure consisting of an ubiquitin-like domain (Ubl), ubiquitin-associated domain 1 (Uba1), XPC-binding domain and Uba2. The protein encoded by this gene plays an important role in nucleotide excision repair and also in delivery of polyubiquitinated proteins to the proteasome. Alternative splicing results in multiple transcript variants encoding multiple isoforms. [provided by RefSeq, Jun 2012],</p>
Function :	<p>domain:The ubiquitin-like domain mediates interaction with MJD.,function:Involved in postreplication repair of UV-damaged DNA. Postreplication repair functions in gap-filling of a daughter strand on replication of damaged DNA .,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the RAD23 family.,similarity:Contains 1 ubiquitin-like domain.,similarity:Contains 2 UBA domains.,subunit:Interacts with MJD. Interacts with HIV-1 Vpr.,</p>
Subcellular Location :	Nucleus.
Expression :	Brain,Pancreas,Testis,
Sort :	13749
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Western blot analysis of lysates from LOVO cells, using RAD23A Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).