


Rad54 Polyclonal Antibody


Catalog No :	YT3970
Reactivity :	Human;Mouse
Applications :	WB;IHC;IF;ELISA
Target :	Rad54
Fields :	>>Homologous recombination
Gene Name :	RAD54L
Protein Name :	DNA repair and recombination protein RAD54-like
Human Gene Id :	8438
Human Swiss Prot No :	Q92698
Mouse Gene Id :	19366
Mouse Swiss Prot No :	P70270
Immunogen :	The antiserum was produced against synthesized peptide derived from human RAD54L. AA range:221-270
Specificity :	Rad54 Polyclonal Antibody detects endogenous levels of Rad54 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:5000. Not yet tested in other applications.
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml

Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	85kD
Cell Pathway :	Homologous recombination;
Background :	<p>The protein encoded by this gene belongs to the DEAD-like helicase superfamily, and shares similarity with <i>Saccharomyces cerevisiae</i> Rad54, a protein known to be involved in the homologous recombination and repair of DNA. This protein has been shown to play a role in homologous recombination related repair of DNA double-strand breaks. The binding of this protein to double-strand DNA induces a DNA topological change, which is thought to facilitate homologous DNA pairing, and stimulate DNA recombination. Alternative splicing results in multiple transcript variants encoding the same protein.[provided by RefSeq, Dec 2008],</p>
Function :	<p>disease:Defects in RAD54L may be a cause of tumor formation. Although alterations are found in a small fraction of primary tumors, these findings provide new insight into genetic instability underlying carcinogenesis.,function:Involved in DNA repair and mitotic recombination. Functions in the recombinational DNA repair (RAD52) pathway. Dissociates RAD51 from nucleoprotein filaments formed on dsDNA. Could be involved in the turnover of RAD51 protein-dsDNA filaments (By similarity). May play also an essential role in telomere length maintenance and telomere capping in mammalian cells.,induction:Expression increases approximately 3-fold in late G1 phase compared to other phases of the cell cycle.,similarity:Belongs to the SNF2/RAD54 helicase family.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,subunit:Interacts with RAD51 through the N</p>
Subcellular Location :	Nucleus .
Expression :	Testis,Thymus,
Tag :	orthogonal
Sort :	13761
No4 :	1
Host :	Rabbit
Modifications :	Unmodified

Products Images



Western blot analysis of lysates from HeLa cells, Jurkat cells, and 293 cells, using RAD54 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).