

PKC θ Polyclonal Antibody

Catalog No :	YT3767
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	PKC θ
Fields :	>>NF-kappa B signaling pathway;>>Autophagy - animal;>>Vascular smooth muscle contraction;>>Th1 and Th2 cell differentiation;>>Th17 cell differentiation;>>T cell receptor signaling pathway;>>Inflammatory mediator regulation of TRP channels;>>Adipocytokine signaling pathway;>>Insulin resistance;>>Shigellosis;>>PD-L1 expression and PD-1 checkpoint pathway in cancer
Gene Name :	PRKCQ
Protein Name :	Protein kinase C theta type
Human Gene Id :	5588
Human Swiss Prot No :	Q04759
Mouse Gene Id :	18761
Mouse Swiss Prot No :	Q02111
Rat Swiss Prot No :	Q9WTQ0
Immunogen :	The antiserum was produced against synthesized peptide derived from human PKC thet. AA range:504-553
Specificity :	PKC θ Polyclonal Antibody detects endogenous levels of PKC θ protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG

Dilution :	<u>WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000.. IF 1:50-200</u>
Purification :	<u>The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.</u>
Concentration :	<u>1 mg/ml</u>
Storage Stability :	<u>-15 °C to -25 °C/1 year(Do not lower than -25 °C)</u>
Observed Band :	<u>78kD</u>
Cell Pathway :	<u>Regulation_Microtubule; Regulation of Actin Dynamics; Stem cell pathway; Insulin Receptor; NF_kappaB; B Cell Receptor; AMPK</u>
Background :	<u>Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role. The protein encoded by this gene is one of the PKC family members. It is a calcium-independent and phospholipid-dependent protein kinase. This kinase is important for T-cell activation. It is required for the activation of the transcription factors NF-kappaB and AP-1, and may link the T cell receptor (TCR) signaling complex to the activation of the transcription factors. [provided by RefSeq, Jul 2008],</u>
Function :	<u>catalytic activity:ATP + a protein = ADP + a phosphoprotein.,cofactor:Magnesium.,domain:The C1 domain, containing the phorbol ester/DAG-type region 1 (C1A) and 2 (C1B), is the diacylglycerol sensor and the C2 domain is a non-calcium binding domain.,enzyme regulation:Three specific sites; Thr-538 (activation loop of the kinase domain), Ser-676 (turn motif) and Ser-695 (hydrophobic region), need to be phosphorylated for its full activation.,function:PKC is activated by diacylglycerol which in turn phosphorylates a range of cellular proteins. PKC also serves as the receptor for phorbol esters, a class of tumor promoters.,function:This is a calcium-independent, phospholipid-dependent, serine- and threonine-specific enzyme. Essential for T-cell receptor (TCR)-mediated T-cell activation, but is dispensable during TCR-dependent thymocyte development. Links the TCR signaling complex to the activ</u>
Subcellular Location :	<u>Cytoplasm. Cell membrane; Peripheral membrane protein. In resting T-cells, mostly localized in cytoplasm. In response to TCR stimulation, associates with lipid rafts and then localizes in the immunological synapse.</u>
Expression :	<u>Expressed in skeletal muscle, T-cells, megakaryoblastic cells and platelets.</u>


Sort : 12766

No4 : 1


Host : Rabbit

Modifications : Unmodified


Products Images



Western Blot analysis of 293 cells using PKC θ Polyclonal Antibody



Immunohistochemistry analysis of paraffin-embedded human brain carcinoma tissue, using PKC θ Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from 293 cells, using PKC θ Antibody. The lane on the right is blocked with the synthesized peptide.