


SAE1 Polyclonal Antibody


Catalog No :	YT4207
Reactivity :	Human;Mouse;Rat
Applications :	WB;IHC;IF;ELISA
Target :	SAE1
Fields :	>>Ubiquitin mediated proteolysis
Gene Name :	SAE1
Protein Name :	SUMO-activating enzyme subunit 1
Human Gene Id :	10055
Human Swiss Prot No :	Q9UBE0
Mouse Gene Id :	56459
Mouse Swiss Prot No :	Q9R1T2
Rat Gene Id :	308384
Rat Swiss Prot No :	Q6AXQ0
Immunogen :	The antiserum was produced against synthesized peptide derived from human AOS1. AA range:221-270
Specificity :	SAE1 Polyclonal Antibody detects endogenous levels of SAE1 protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500 - 1:2000. IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.

Purification :	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	38kD
Cell Pathway :	Ubiquitin mediated proteolysis;
Background :	SUMO1 activating enzyme subunit 1(SAE1) Homo sapiens Posttranslational modification of proteins by the addition of the small protein SUMO (see SUMO1; MIM 601912), or sumoylation, regulates protein structure and intracellular localization. SAE1 and UBA2 (MIM 613295) form a heterodimer that functions as a SUMO-activating enzyme for the sumoylation of proteins (Okuma et al., 1999 [PubMed 9920803]).[supplied by OMIM, Mar 2010],
Function :	function:The dimeric enzyme acts as a E1 ligase for SUMO1, SUMO2, SUMO3, and probably SUMO4. It mediates ATP-dependent activation of SUMO proteins and formation of a thioester with a conserved cysteine residue on SAE2.,pathway:Protein modification; protein sumoylation.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the ubiquitin-activating E1 family.,subunit:Heterodimer of SAE1 and SAE2. The complex binds SUMO proteins via SAE2.,tissue specificity:Expression level increases during S phase and drops in G2 phase (at protein level),,
Subcellular Location :	Nucleus .
Expression :	Expression level increases during S phase and drops in G2 phase (at protein level).
Sort :	14742
No4 :	1
Host :	Rabbit
Modifications :	Unmodified


Products Images




Western Blot analysis of various cells using SAE1 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Immunofluorescence analysis of HUVEC cells, using AOS1 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from 293 cells, using AOS1 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).