


Raftlin Polyclonal Antibody


Catalog No :	YT3989
Reactivity :	Human;Mouse
Applications :	WB;IHC
Target :	Raftlin
Gene Name :	RFTN1
Protein Name :	Raftlin
Human Gene Id :	23180
Human Swiss Prot No :	Q14699
Mouse Swiss Prot	Q6A0D4
Immunogen :	The antiserum was produced against synthesized peptide derived from human RFTN1. AA range:432-481
Specificity :	Raftlin Polyclonal Antibody detects endogenous levels of Raftlin protein.
Formulation :	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source :	Polyclonal, Rabbit,IgG
Dilution :	WB 1:500-2000;IHC 1:50-300
Purification :	The antibody was affinity-purified from rabbit antiserum by affinity- chromatography using epitope-specific immunogen.
Concentration :	1 mg/ml
Storage Stability :	-15°C to -25°C/1 year(Do not lower than -25°C)
Observed Band :	65kD



Background :	function:May play a pivotal role in the formation and/or maintenance of lipid rafts. May regulate B-cell antigen receptor-mediated signaling.,similarity:Belongs to the raftlin family.,subcellular location:Associates with lipid rafts.,tissue specificity:Specifically expressed by B-cells (at protein level).,
Function :	function:May play a pivotal role in the formation and/or maintenance of lipid rafts. May regulate B-cell antigen receptor-mediated signaling.,similarity:Belongs to the raftlin family.,subcellular location:Associates with lipid rafts.,tissue specificity:Specifically expressed by B-cells (at protein level).,
Subcellular Location :	Cell membrane ; Lipid-anchor . Cytoplasm . Membrane raft . Endosome . Early endosome . Translocates from cytoplasm to cell membrane where it colocalizes with poly (I:C) and then moves to endosomes where it colocalizes with TLR3 (PubMed:21266579). Translocates from cytoplasm to cell membrane where it colocalizes with TLR4 and then together with TLR4 moves to endosomes, upon lipopolysaccharide stimulation (PubMed:27022195)
Expression :	Expressed in B-cells (at protein level) (PubMed:12805216). Expressed in dendritic cells and macrophages (PubMed:27022195, PubMed:21266579).
Sort :	13797
No4 :	1
Host :	Rabbit
Modifications :	Unmodified







RFTN1

117-

85-

48-


34-

26-

19-

Western blot analysis of RFTN1 Antibody. The lane on the right is blocked with the RFTN1 peptide.

Western blot analysis of the lysates from RAW264.7cells using RFTN1 antibody.



Immunohistochemical analysis of paraffin-embedded human uterus. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).