

Ribosomal Protein LP2 Polyclonal Antibody

YT4120 **Catalog No:**

Human; Mouse; Rat Reactivity:

Applications: IHC;IF;ELISA

Target: RPLP2

Fields: >>Ribosome;>>Coronavirus disease - COVID-19

Gene Name: RPLP2

Protein Name: 60S acidic ribosomal protein P2

P99027

Human Gene Id: 6181

Human Swiss Prot

P05387

No:

Mouse Gene Id: 67186

Mouse Swiss Prot

No:

1.00912e+008 Rat Gene Id:

Rat Swiss Prot No: P02401

The antiserum was produced against synthesized peptide derived from human Immunogen:

RPLP2. AA range:21-70

Ribosomal Protein LP2 Polyclonal Antibody detects endogenous levels of **Specificity:**

Ribosomal Protein LP2 protein.

Formulation: Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: IHC 1:100 - 1:300. ELISA: 1:40000.. IF 1:50-200

1/3



Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Molecularweight: 12kD

Cell Pathway: Ribosome;

Background : Ribosomes, the organelles that catalyze protein synthesis, consist of a small

40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. This gene encodes a ribosomal phosphoprotein that is a component of the 60S subunit. The protein, which is a functional equivalent of the E. coli L7/L12 ribosomal protein, belongs to the L12P family of ribosomal proteins. It plays an important role in the elongation step of protein synthesis. Unlike most ribosomal proteins, which are basic, the encoded protein is acidic. Its C-terminal end is nearly identical to the C-terminal ends of the ribosomal phosphoproteins P0 and P1. The P2 protein can interact with P0 and P1 to form a pentameric complex consisting of P1 and P2 dimers, and a P0 monomer. The protein is located in the cytoplasm. As is typical

for genes

Function: function:Plays an important role in the elongation step of protein

synthesis., similarity: Belongs to the ribosomal protein L12P family., subunit: P1 and

P2 exist as dimers at the large ribosomal subunit.,

Subcellular Location:

cytosol, ribosome, focal adhesion, membrane, cytosolic large ribosomal subunit, preribosome, large subunit precursor, extracellular exosome,

Expression: Epithelium, Kidney, Liver, Mammary carcinoma, Ovary, Pituitary, P


Sort: 14487

No4: 1

Host: Rabbit

Modifications: Unmodified

Products Images



Immunohistochemistry analysis of paraffin-embedded human prostate carcinoma tissue, using RPLP2 Antibody. The picture on the right is blocked with the synthesized peptide.