

Septin 1 Polyclonal Antibody

Catalog No: YT4245

Reactivity: Human; Mouse; Rat

Applications: WB;IHC;IF;ELISA

Target: Septin 1

Fields: >>Bacterial invasion of epithelial cells

Gene Name: SEPT1

Protein Name: Septin-1

Human Gene Id: 1731

Human Swiss Prot

Q8WYJ6

P42209

No:

Mouse Gene ld: 54204

Mouse Swiss Prot

No:

Rat Gene Id: 293507

Rat Swiss Prot No: Q5EB96

Immunogen: The antiserum was produced against synthesized peptide derived from human

SEPT1. AA range:181-230

Specificity: Septin 1 Polyclonal Antibody detects endogenous levels of Septin 1 protein.

Formulation : Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

Source: Polyclonal, Rabbit, IgG

Dilution: WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000.. IF 1:50-200

1/3



Purification: The antibody was affinity-purified from rabbit antiserum by affinity-

chromatography using epitope-specific immunogen.

Concentration: 1 mg/ml

Storage Stability: -15°C to -25°C/1 year(Do not lower than -25°C)

Observed Band: 41kD

Background: septin 1(SEPT1) Homo sapiens This gene is a member of the septin family of

GTPases. Members of this family are required for cytokinesis and the

maintenance of cellular morphology. This gene encodes a protein that can form homo- and heterooligomeric filaments, and may contribute to the formation of neurofibrillary tangles in Alzheimer's disease. Alternatively spliced

transcript variants have been found but the full-length nature of these variants has

not been determined. [provided by RefSeq, Dec 2012],

Function: function:Involved in cytokinesis .,similarity:Belongs to the septin

family., subunit: May assemble into a multicomponent structure.,

Subcellular Cytoplasm . Cytoplasm, cytoskeleton . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Midbody. Remains at the centrosomes and the

organizing center, centrosome. Midbody. Remains at the centrosomes and the nearby microtubules throughout mitosis. Localizes to the midbody during

cytokinesis.

Expression: Expressed at high levels in lymphoid and hematopoietic tissues.

Sort: 16232


No4: 1

Host: Rabbit


Modifications: Unmodified

Products Images


2/3



Western Blot analysis of various cells using Septin 1 Polyclonal Antibody



Western blot analysis of lysates from Jurkat cells, using SEPT1 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded human spleen tissue. 1,primary Antibody was diluted at 1:200(4° overnight). 2, Sodium citrate pH 6.0 was used for antigen retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200